

Baryon fraction from the BAO amplitude: A consistent approach to parametrizing perturbation growth

Andrea Crespi,^{1,2,*} Will J. Percival,^{1,2,3} Alex Krolewski,^{1,2} Marco Bonici,^{1,2} Hanyu Zhang,^{1,2} J. Aguilar,⁴ S. Ahlen,⁵
 A. Anand,⁴ D. Bianchi,^{6,7} D. Brooks,⁸ E. Chaussidon,⁴ T. Claybaugh,⁴ A. Cuceu,⁴ A. de la Macorra,⁹ P. Doel,⁸
 S. Ferraro,^{4,10} A. Font-Ribera,¹¹ J. E. Forero-Romero,^{12,13} E. Gaztañaga,^{14,15,16} G. Gutierrez,¹⁷ J. Guy,⁴
 H. K. Herrera-Alcantar,^{18,19} D. Huterer,^{20,21} M. Ishak,²² D. Joyce,²³ D. Kirkby,²⁴ T. Kisner,⁴ A. Kremin,⁴ O. Lahav,⁸
 C. Lamman,²⁵ M. Landriau,⁴ L. Le Guillou,²⁶ M. E. Levi,⁴ M. Manera,^{27,28} P. Martini,^{25,29,30} A. Meisner,²³
 R. Miquel,^{27,31} S. Nadathur,¹⁵ N. Palanque-Delabrouille,^{4,19} C. Poppett,^{4,10,32} F. Prada,³³ I. Pérez-Ràfols,³⁴ G. Rossi,³⁵
 L. Samushia,^{36,37,38} E. Sanchez,³⁹ D. Schlegel,⁴ M. Schubnell,^{20,21} H. Seo,⁴⁰ J. H. Silber,⁴ D. Sprayberry,²³ G. Tarlé,²¹
 B. A. Weaver,²³ R. Zhou,⁴ and H. Zou⁴¹

¹Waterloo Centre for Astrophysics, *University of Waterloo*, Waterloo, Ontario N2L 3G1, Canada

²Department of Physics and Astronomy, *University of Waterloo*, Waterloo, Ontario N2L 3G1, Canada

³Perimeter Institute for Theoretical Physics, 31 Caroline St. North, Waterloo, Ontario N2L 2Y5, Canada

⁴Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, California 94720, USA

⁵Department of Physics, *Boston University*, 590 Commonwealth Avenue, Boston, Massachusetts 02215 USA

⁶Dipartimento di Fisica “Aldo Pontremoli,” *Università degli Studi di Milano*, Via Celoria 16, I-20133 Milano, Italy

⁷INAF-Osservatorio Astronomico di Brera, Via Brera 28, 20122 Milano, Italy

⁸Department of Physics and Astronomy, *University College London*, Gower Street, London, WC1E 6BT, United Kingdom

⁹Instituto de Física, *Universidad Nacional Autónoma de México*, Circuito de la Investigación Científica, Ciudad Universitaria, Cd. de México C. P. 04510, México

¹⁰*University of California*, Berkeley, 110 Sproul Hall #5800 Berkeley, California 94720, USA

¹¹Institut de Física d’Altes Energies (IFAE), *The Barcelona Institute of Science and Technology*, Edifici Cn, Campus UAB, 08193, Bellaterra Barcelona, Spain

¹²Departamento de Física, *Universidad de los Andes*, Cra. 1 No. 18A-10, Edificio Ip, CP 111711, Bogotá, Colombia

¹³Observatorio Astronómico, *Universidad de los Andes*, Cra. 1 No. 18A-10, Edificio H, CP 111711 Bogotá, Colombia

¹⁴Institut d’Estudis Espacials de Catalunya (IEEC), c/ Esteve Terradas 1, Edifici RDIT, Campus PMT-UPC, 08860 Castelldefels, Spain

¹⁵Institute of Cosmology and Gravitation, *University of Portsmouth*, Dennis Sciana Building, Portsmouth, PO1 3FX, United Kingdom

¹⁶*Institute of Space Sciences*, ICE-CSIC, Campus UAB, Carrer de Can Magrans s/n, 08913 Bellaterra, Barcelona, Spain

¹⁷*Fermi National Accelerator Laboratory*, PO Box 500, Batavia, Illinois 60510, USA

¹⁸Institut d’Astrophysique de Paris. 98 bis boulevard Arago. 75014 Paris, France

¹⁹IRFU, CEA, *Université Paris-Saclay*, F-91191 Gif-sur-Yvette, France

²⁰Department of Physics, *University of Michigan*, 450 Church Street, Ann Arbor, Michigan 48109, USA

²¹*University of Michigan*, 500 S. State Street, Ann Arbor, Michigan 48109, USA

²²Department of Physics, *The University of Texas at Dallas*, 800 W. Campbell Road, Richardson, Texas 75080, USA

²³NSF NOIRLab, 950 N. Cherry Avenue, Tucson, Arizona 85719, USA

²⁴Department of Physics and Astronomy, *University of California*, Irvine, California 92697, USA

²⁵*The Ohio State University*, Columbus, 43210 Ohio, USA

²⁶Sorbonne Université, CNRS/IN2P3, *Laboratoire de Physique Nucléaire et de Hautes Energies (LPNHE)*, FR-75005 Paris, France

²⁷Institut de Física d’Altes Energies (IFAE), *The Barcelona Institute of Science and Technology*, Edifici Cn, Campus UAB, 08193, Bellaterra (Barcelona), Spain

²⁸Departament de Física, Serra Húnter, *Universitat Autònoma de Barcelona*, 08193 Bellaterra (Barcelona), Spain

²⁹Center for Cosmology and AstroParticle Physics, *The Ohio State University*, 191 West Woodruff Avenue, Columbus, Ohio 43210, USA

³⁰Department of Astronomy, *The Ohio State University*, 4055 McPherson Laboratory, 140 W 18th Avenue, Columbus, Ohio 43210, USA

³¹*Institució Catalana de Recerca i Estudis Avançats, Passeig de Lluís Companys, 23, 08010 Barcelona, Spain*

³²*Space Sciences Laboratory, University of California, Berkeley, 7 Gauss Way, Berkeley, California 94720, USA*

³³*Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía, s/n, E-18008 Granada, Spain*

³⁴*Departament de Física, EEBE, Universitat Politècnica de Catalunya, c/Eduard Maristany 10, 08930 Barcelona, Spain*

³⁵*Department of Physics and Astronomy, Sejong University, 209 Neungdong-ro, Gwangjin-gu, Seoul 05006, Republic of Korea*

³⁶*Abastumani Astrophysical Observatory, Tbilisi, GE-0179, Georgia*

³⁷*Department of Physics, Kansas State University, 116 Cardwell Hall, Manhattan, Kansas 66506, USA*

³⁸*Faculty of Natural Sciences and Medicine, Ilia State University, 0194 Tbilisi, Georgia*

³⁹*CIEMAT, Avenida Complutense 40, E-28040 Madrid, Spain*

⁴⁰*Department of Physics and Astronomy, Ohio University, 139 University Terrace, Athens, Ohio 45701, USA*

⁴¹*National Astronomical Observatories, Chinese Academy of Sciences, A20 Datun Road, Chaoyang District, Beijing, 100101, People's Republic of China*



(Received 19 December 2025; revised 9 June 2026; accepted 24 July 2026; published 25 August 2026)

Galaxy clustering constrains the baryon fraction Ω_b/Ω_m through the amplitude of baryon acoustic oscillations and the suppression of perturbations entering the horizon before recombination. This produces a different prerecombination distribution of baryons and dark matter. After recombination, the gravitational potential responds to both components in proportion to their mass, allowing robust measurement of the baryon fraction. This is independent of new-physics scenarios altering the recombination background (e.g. early dark energy). The accuracy of such measurements does, however, depend on how baryons and cold dark matter (CDM) are modeled in the power spectrum. Previous template-based splitting relied on approximate transfer functions that neglected part of information. We present a new method that embeds an extra parameter controlling the balance between baryons and dark matter in the growth terms of the perturbation equations in the CAMB Boltzmann solver. This approach captures the baryonic suppression of CDM prior to recombination, avoids inconsistencies, and yields a clean parametrization of the baryon fraction in the linear power spectrum, separating out the simple physics of growth due to the combined matter potential. We implement this framework in an analysis pipeline using effective field theory of large-scale structure with halo occupation distribution-informed priors and validate it against noiseless Λ CDM and early dark energy cosmologies with Dark Energy Spectroscopic Instrument (DESI)-like errors. The new scheme achieves comparable precision to previous splitting while reducing systematic biases, providing a more robust way to baryon-fraction measurements. In combination with big bang nucleosynthesis constraints on the baryon density and Alcock-Paczyński estimates of the matter density, these results strengthen the use of baryon fraction measurements to derive a Hubble constant from energy densities, with future DESI and Euclid data expected to deliver competitive constraints.

DOI: 10.1103/fcb6-7xpx

I. INTRODUCTION

The flat Λ CDM model parameters have been constrained to percent level precision by observations of the cosmic microwave background (CMB) [1], which provides strong evidence for this model. However, tensions with other data have shifted the key questions from those of parameter estimation to consistency tests. One of the strongest tensions is the $\sim 4\text{--}6\sigma$ tension between local distance-ladder determinations of the Hubble constant, $H_0 = 73.04 \pm 1.04 \text{ km s}^{-1} \text{ Mpc}^{-1}$, combining Cepheids

and supernovae [2], and the CMB-inferred value, $H_0 = 67.37 \pm 0.54 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [1]. Similarly, baryon acoustic oscillation (BAO) measurements yield a comparable low value when combined with the standard ruler of the sound horizon, calibrated using big bang nucleosynthesis (BBN) data [3]. Explanations for the tension range from unidentified systematic errors in the datasets to extensions of Λ CDM that invoke new physics—such as early dark energy (EDE) [4–7] (see also [8,9] and [10] for other early time modifications that can potentially resolve the Hubble tension). Although the most recent ACT DR6 results [11] alone are compatible with a nonzero EDE component, the preference for EDE disappears once supernova data and

*Contact author: a2crespi@uwaterloo.ca

DESI DR2 BAO measurements [12] are included in the analysis. Therefore, independent and complementary measurements of H_0 are essential to understand the origin of this discrepancy.

Recently, several methods have been developed to determine H_0 independently on the sound horizon and recombination physics. Philcox *et al.* [13] (using methods developed in [14–16]) infer H_0 from the matter-radiation equality scale using Baryon Oscillation Spectroscopic Survey (BOSS) full-shape power spectra and CMB lensing, marginalizing over the sound horizon to remove acoustic information. Bahr-Kalus *et al.* [17] directly detect the turnover in the matter power spectrum and use it as a standard ruler, yielding a sound-horizon-free estimate of the Hubble constant. Zaborowski *et al.* [18] extend this strategy by analyzing the full-shape clustering of multiple tracers, rescaling the power spectrum to explicitly marginalize over the sound horizon, and combining with CMB lensing and uncalibrated Supernovae (SNe) to reach sub-3% precision. In all cases, the calibration depends on equality physics or direct shape information rather than on the sound horizon at drag epoch, making the resulting H_0 measurements independent of recombination-era acoustic physics. In a recent paper [19], Krolewski *et al.* presented a new method for inferring the Hubble constant that avoids using CMB power-spectrum data and does not require calibrating the BAO with a predetermined sound-horizon scale. This method combines three independent pieces of information. First, a measurement of the matter density in units of the critical density, Ω_m is obtained from the Alcock-Paczyński (AP) effect applied to large-scale structure, which provides a purely geometric constraint based on anisotropic distortions of BAO (or cosmic voids [20]). Second, the physical baryon density $\Omega_b h^2$ is constrained by BBN. The remaining ingredient required to determine H_0 is the baryon fraction, $f_b = \Omega_b / \Omega_m$. The three measurements together constrain the critical density, which in turn depends on the Hubble constant. In their companion work [21] (hereafter KP25), Krolewski and Percival also introduced a new method to infer f_b from the amplitude of the baryonic signal in the galaxy power spectrum. This baryonic signature consists of two effects: the acoustic oscillations imprinted in the range $k \sim 0.02\text{--}0.3 \text{ h Mpc}^{-1}$ and a scale-dependent suppression on subsound-horizon scales. In particular, Fig. 1 in KP25 shows that the BAO amplitude can recover the true baryon fraction f_b , independent of whether the underlying cosmology follows Λ CDM or an EDE model. Thus, it provides the key ingredient for a energy density based measurement of H_0 . In this work, we develop a more robust method for measuring the baryon fraction than that adopted by KP25, by introducing an additional free parameter to separate the growth contributions of baryons and cold dark matter in the perturbation equations. We begin by contextualizing the problem in Sec. II, present the new method in Sec. III, and describe how it can be integrated into a pipeline for

measuring H_0 in Sec. IV. In Sec. V, we provide a detailed comparison between the original KP25 template method and our perturbation-level implementation on noiseless theory vectors spanning Λ CDM and EDE cosmologies, and our conclusions are summarized in Sec. VI. In this work, we focus only on the validation of the method on noiseless data vectors; the application to the Dark Energy Spectroscopic Instrument (DESI; [22]), Data Release 1 (DR1; [23]) sample is provided in our companion paper Krolewski *et al.* (in prep).

II. MEASURING THE BARYON FRACTION FROM GROWTH

Within galaxy redshift surveys, the imprint left by baryons is one of the most robust physical signatures that can be extracted. Classical BAO analyses (e.g. [24]) have focused on measuring the observed positions of the feature along and across the line of sight, which scale as D_H/r_d and D_M/r_d , with $D_H(z) \equiv c/H(z)$, $D_M(z) \equiv (1+z)D_A(z)$, where $H(z)$ is the Hubble flow, $D_A(z)$ is the angular diameter distance, and r_d the comoving sound horizon at the drag epoch. The sound horizon depends nontrivially on Ω_b , Ω_m , and h , requiring a complex physical modeling. Nevertheless, it is predicted to very high precision once the cosmology is fixed. In linear theory, the cosmic evolution of density perturbations is encapsulated in scale-dependent transfer functions: each component—baryons, cold dark matter, photons, neutrinos—has its own $T_i(k)$ that translates primordial fluctuations into present-day amplitudes. For an energy-density dominated by matter, the overall transfer function is

$$T(k) = \frac{\Omega_b}{\Omega_{bc}} T_b(k) + \frac{\Omega_c}{\Omega_{bc}} T_c(k), \quad (1)$$

where $\Omega_{bc} \equiv \Omega_b + \Omega_c$, with the baryonic and cold-dark-matter components entering in proportion to their energy-density fractions. This function, feeds directly into the linear clustering power spectrum

$$P(k, z) = A_s \left(\frac{kh}{0.05} \right)^{n_s-1} 2\pi^2 k h^4 T^2(k) D^2(z), \quad (2)$$

where A_s is the amplitude of primordial power spectrum, n_s is the scalar spectral index, h is the dimensionless Hubble parameter, $T(k)$ is the matter transfer function, and $D(z)$ is the linear growth factor. Equation (1) exhibits an explicit and simple dependence on the fraction Ω_b/Ω_{bc} .

To isolate the information on the baryon fraction coming from the growth of perturbations, we can consider replacing the weighting between T_b and T_c with a free parameter (hereafter γ_b), such that the weights for the two terms are γ_b and $(1 - \gamma_b) * \gamma_b$ is an estimator of the ratio between baryon and baryon + CDM densities coming from just the growth of perturbations.

Before photon decoupling, the baryon and CDM transfer functions differ significantly. The CDM component is a smooth function of k , while the baryonic component exhibits acoustic oscillations imprinted by the photon-baryon fluid. On scales well below the sound horizon, the baryon transfer function can be approximated as an oscillatory term modulated by a damping envelope, reflecting silk damping and the suppression of baryon perturbations prior to recombination. After the drag epoch, the baryonic and CDM perturbations grow together under the summed gravitational potential. As a result, the shape of the total transfer function is fixed and subsequent evolution only changes the overall amplitude of the perturbations in a scale-independent way. Since f_b is constant with redshift in standard cosmological models such as Λ CDM and the EDE scenarios considered in this work, $T_b(k)$ and $T_c(k)$ can be evaluated at any convenient epoch (rescaling their amplitudes for the correct redshift), and recombined as in Eq. (1). This freedom can be used to potentially maximize the contrast between the two transfer functions and improve the precision of the γ_b measurement. However, as explained in KP25, the two transfer functions must be chosen with care. At low redshift ($z = 0$), the two components have already evolved together and therefore look very similar, removing the information needed to distinguish their relative contributions. At the opposite extreme, defining the transfer functions at recombination maximizes their differences, but does not reflect the final state of the acoustic oscillations, which continue to propagate after decoupling. Intermediate redshifts partially capture this evolution, but still mix residual baryonic oscillations with the emerging CDM response, making the separation less clean. To overcome these issues, KP25 adopted a hybrid approach. The CDM transfer function was taken from the Eisenstein and Hu (EH) fitting formula [25], which provides a smooth, nonoscillatory description of the CDM component. The baryonic contribution was then defined as the difference between the total matter transfer function from CLASS [26] at $z = 0$ and the EH CDM component, rescaled by the baryon fraction

$$T_b(k) = \left(T_{cb}(k, z=0) - \frac{\Omega_c}{\Omega_{cb}} T_c^{\text{EH}}(k) \right) \frac{\Omega_{cb}}{\Omega_b}. \quad (3)$$

This construction allows one to retain the final position of the acoustic oscillations while maximizing the separation between baryonic and CDM contributions. However, as we are going to further explore in Sec. III, this splitting choice has two major limitations. First, the numerical precision of EH transfer functions puts a bound on the precision of γ_b measurements. Second, and more importantly, this approach fixes the shape of T_c as it is at the drag epoch, neglecting the effect of baryons on the growth of CDM perturbations prior to recombination. This omission reduces the physical self-consistency and discards part of the physical information carried by a full perturbation evolution.

For this analysis, we assumed a single set of bias parameters when comparing the galaxy field to both the CDM and baryon density fields. However, at early times, baryons and CDM potentially have different velocities and densities. This relative evolution can induce additional bias contributions, introducing two extra terms, $b_{\delta_{bc}}$ and $b_{v_{bc}^2}$, that are generally scale dependent. Similarly, compensated isocurvature perturbations (CIPs) [27–29] correspond to primordial fluctuations in the baryon density that are compensated by opposite fluctuations in the CDM density, leaving the total matter density unchanged. These perturbations can also generate scale-dependent bias contributions. Although these effects can, in principle, be modeled, they are expected to be negligible at the level of precision of current data. A more detailed treatment of these contributions is therefore deferred to future work. A qualitative discussion of the impact of relative velocity effects can be found in Appendix A of the companion paper [30].

III. MODELING THE GROWTH THROUGH PERTURBATION EVOLUTION

We propose a new approach that embeds γ_b directly in the linear perturbation equations of the Boltzmann solver CAMB [31]. The baryon and CDM densities are rescaled according to the new parameter, so the full temporal evolution of both fluids responds coherently to the chosen γ_b . The thermal history and the adiabatic initial conditions, governed by the background cosmological parameters, are unaltered. This means that the new parameter is able to extract the dependence of perturbation growth only on the baryon fraction. This type of approach is not new. Several works in the literature have split parameters into two distinct counterparts, each responsible for different physical phenomena, motivated by different goals. For example, the separation between a total matter energy density parameter for growth and one for geometry to test the Λ CDM model in [32], or the splitting of the baryon density into a parameter controlling the ionization history and another governing the rest of the baryon effects in [33]. There are different ways to separate the dependence on the growth in the physical processes and in their equations, each encapsulating distinct physical aspects of the evolution of perturbations. We consider a number of options to better explain why our chosen method is the most coherent and complete after first reviewing the physical equations governing the evolutions of perturbations.

A. Evolution of perturbations

CAMB¹ numerically integrates the linearized Einstein-Boltzmann system in synchronous gauge, evolving all

¹Particularly valuable for understanding the equations for the evolution of the perturbations and their numerical implementation were Prof. Antony M. Lewis's PhD thesis, *Geometric Algebra and Covariant Methods in Physics and Cosmology* [34], as well as his CAMB NOTES [35].

species in conformal time τ so that the gravitational potentials sourced by one component instantaneously feed back on every other. The background expansion obeys the Friedmann equation

$$\mathcal{H}^2 \equiv \left(\frac{a'}{a}\right)^2 = \frac{8\pi G}{3} a^2 \sum_i \bar{\rho}_i(\tau), \quad (4)$$

where a prime denotes $d/d\tau$ and the sum runs over photons, massless and massive neutrinos, cold dark matter, baryons, and dark energy. Metric perturbations are described in synchronous-gauge by the two scalar potentials h and η . The Hamiltonian and momentum equations result in

$$\begin{aligned} k^2 \eta - \frac{1}{2} \mathcal{H} h' &= 4\pi G a^2 \sum_i \bar{\rho}_i \delta_i \\ k \eta' &= 4\pi G a^2 \sum_i (\bar{\rho}_i + \bar{P}_i) v_i, \end{aligned} \quad (5)$$

with density contrast $\delta_i \equiv \delta\rho_i/\bar{\rho}_i$ and pressure P_i . Cold dark matter evolves according to

$$\delta'_c = -\frac{1}{2} h', \quad v'_c = 0, \quad (6)$$

while baryons are tightly coupled to photons through Thomson scattering before decoupling. This tight coupling results in a high opacity τ_c^{-1} , and in this regime, the full Boltzmann hierarchy for photons becomes stiff and extremely inefficient. To address this, CAMB implements a tight-coupling approximation by expanding in τ_c , valid when $\epsilon \equiv \max(k\tau_c, \mathcal{H}\tau_c) \ll 1$:

$$\begin{aligned} \delta'_b &= -\frac{1}{2} h' - k v_b \\ v'_b &= \frac{1}{1+R} \left[k c_s^2 \delta_b + \frac{k}{4} R (\delta_\gamma - 2\beta_2 \pi_\gamma) - \mathcal{H} v_b \right. \\ &\quad + \left(\frac{2\mathcal{H}}{1+R} + (\ln \tau_c^{-1})' \right) \frac{3R}{4(1+R)} \Delta \\ &\quad - \frac{R\tau_c}{4(1+R)^2} (4(\mathcal{H}' + \mathcal{H}^2) v_b \\ &\quad \left. + k(2\mathcal{H}\delta_\gamma + \delta'_\gamma - 4c_s^2 \delta'_b) \right]. \end{aligned} \quad (7)$$

Here τ_c is the optical depth, $R \equiv 4\bar{\rho}_\gamma/3\bar{\rho}_b$, and c_s is the baryon sound speed. The leading term $k c_s^2 \delta_b$ represents the baryon pressure term, while $\frac{R}{4} (\delta_\gamma - 2\beta_2 \pi_\gamma)$ encodes the momentum input from photon overdensities δ_γ and the anisotropic stress π_γ , which contributes to quadrupole radiation pressure and is computed from the baryon velocity and the metric shear $\sigma = (h' + 6\eta'_s)/(2k)$. The coefficient β_2 accounts for the polarization correction to the quadrupole moment ($\beta_2 = 1$ in the absence of

polarization). The term $-\mathcal{H}v_b$ reflects Hubble drag due to the expansion of the Universe. The Δ term captures the photon-baryon slip, i.e., the relative velocity between photons and baryons. These approximations are necessary to ensure numerical stability and continuity at early times, particularly for small scales where the photon-baryon fluid exhibits acoustic oscillations.

Baryons transition, then, to a epoch of nontight coupling where

$$v'_b \approx -\mathcal{H}v_b + c_s^2 k \delta_b + \frac{k}{4} R \delta_\gamma, \quad (8)$$

which remains valid even after photon decoupling to account for residual baryon-photon interactions from drag epoch onward. The transition between the tight-coupling approximation and the nontight-coupling regime is performed continuously.

Photon evolution is described by the temperature multipoles Θ_ℓ , which satisfy the Boltzmann hierarchy with a baryon-photon scattering term controlled by τ'_c . Massless neutrinos have an analogous system of equations for the multipoles F_ℓ , but without the scattering term τ'_c . Massive neutrinos are integrated in momentum shells with a similar hierarchy weighted by the Fermi-Dirac distribution. Anisotropic stress from free-streaming species feeds back on the metric through Eq. (5), damping potential decay inside the horizon.

Evolving this set of equations all together is essential because each species continuously sources the metric potentials that, in turn, feed back on every other species. Photon-baryon pressure alters the wells into which cold dark matter falls; CDM gravity reshapes the acoustic oscillations of the photon-baryon fluid; free-streaming neutrinos and dark energy likewise modify the same potentials at later times. This tight interdependence means that there is no perfect, redshift-zero transfer function splitting: the baryonic imprint accumulates incrementally at every integration step, inseparable from the concurrent CDM response. Quantifying their individual contributions must therefore compare full, self-consistent integrations rather than isolating them at the end.

B. Modification strategy

We introduce a new parameter, the growth baryon fraction γ_b , allowing this to be different from the cosmological baryon fraction f_b fixed by the background parameters. This allows the growth of perturbations to respond to a different balance of baryonic and dark-matter contributions than is actually present. This is clearly unphysical, but is only used as a mechanism to split the information from the different physical processes. In a fit to data it allows us to extract only the information on the baryon fraction from perturbation growth by considering only the constraint on γ_b . One possible approach is to introduce γ_b only in the

perturbation evolution after decoupling. In this case, all evolution equations are modified at a transition time t_{dec} : before decoupling, they depend on the cosmological baryon fraction, while after decoupling they depend on the new growth baryon fraction, but only in the terms relevant for structure growth. Here, “all evolution equations” refers to the full set of coupled perturbation equations governing all species, including baryons, CDM, photons, and neutrinos, which continuously source the metric perturbations. Because of this coupling, an abrupt modification at t_{dec} introduces a discontinuity in the evolution of the system. This procedure leads to several undesirable features, including step-like variations in the power spectrum. Moreover, part of the information on the baryon-CDM interplay is generated before decoupling and cannot be cleanly separated by switching the equations at a later time. As a result, residual baryonic oscillations persist even in the limit $\gamma_b \rightarrow 0$, while for large γ_b the overall amplitude is affected by prerecombination suppression effects that are not consistently removed. These behaviors reflect the fact that the perturbations have already evolved jointly before the transition and therefore retain a memory of the predecoupling dynamics. A second strategy is to modify the metric perturbations at all times using γ_b , while applying the split in the other evolution equations only after decoupling. In this case, the coupling between species remains partially inconsistent across the transition, since one component is modified continuously while the others are switched abruptly. This leads to a configuration that is even less self-consistent, and similarly produces features in the clustering signal that are difficult to interpret physically.

More generally, we find that abruptly modifying the evolution equations at a specific time introduces artificial features in the final power spectrum that lack a clear physical interpretation. By contrast, when the perturbation equations are modified consistently at all times, as in the approach presented in this work, these features disappear and the resulting evolution remains physically coherent.

C. The new approach

We favor a scenario where we replace the cosmological baryon fraction with its growth counterpart in the equations at all cosmic times, including the prerecombination era. Importantly, we implemented this by carefully considering all the relevant equations in CAMB so that the growth baryon fraction modifies only structure formation; the thermal history of the Universe remains entirely set by the standard cosmological parameters. This approach removes any discontinuity in the dynamics of perturbations and is therefore self-consistent, including baryon induced suppression of CDM perturbations before decoupling. The collapse of matter overdensities, and the subsequent growth of structure, is a delicate interplay between gravitational collapse, expansion of the Universe and mutual interaction between different species. Hence, we have to modify most

of the terms in the equations of Sec. III A. First of all, different baryon and CDM abundances would alter the expansion of the Universe through the Friedmann equation, Eq. (4). Rescaling the baryon and CDM densities with γ_b , but preserving the total matter density, we obtain

$$\mathcal{H}^2 = \frac{8\pi G}{3} a^2 \left(\frac{\gamma_b}{f_b} \bar{\rho}_b(\tau) + \frac{1-\gamma_b}{1-f_b} \bar{\rho}_c(\tau) + \sum_i \bar{\rho}_i(\tau) \right), \quad (9)$$

where the final sum runs over the remaining species. The response of the metric perturbation is also altered in Eq. (5) to

$$\begin{aligned} k^2 \eta - \frac{1}{2} \mathcal{H} h' &= 4\pi G a^2 \left(\frac{\gamma_b}{f_b} \bar{\rho}_b \delta_b + \frac{1-\gamma_b}{1-f_b} \bar{\rho}_c \delta_c + \sum_i \bar{\rho}_i \delta_i \right) \\ k \eta' &= 4\pi G a^2 \left(\frac{\gamma_b}{f_b} \bar{\rho}_b v_b + \sum_i (\bar{\rho}_i + \bar{P}_i) v_i \right), \end{aligned} \quad (10)$$

where, in the second line, there are no CDM terms since in the synchronous gauge it has zero velocity. The equations for δ'_c and δ'_b are not modified further since h' already accounts for the correct growth baryon fraction. Equation (7) for the baryon velocity needs to be treated carefully. The terms that depend on the thermal history of the Universe are left unchanged. This means that the terms responsible for the position of the acoustic peaks do not change, whereas those that control the suppression of baryon growth are modified according to γ_b . With this prescription, the baryon-velocity equation in the tight-coupling regime can be written as

$$\begin{aligned} v'_b &= \frac{1}{1+R} \left[k c_s^2 \delta_b^{(\gamma_b)} + \frac{k}{4} R (\delta_\gamma^{(\gamma_b)} - 2\beta_2 \pi_\gamma^{(\gamma_b)}) - \mathcal{H}^{(\gamma_b)} v_b \right. \\ &\quad + \left(\frac{2\mathcal{H}^{(\gamma_b)}}{1+R} + (\ln \tau_c^{-1})' \right) \frac{3R}{4(1+R)} \Delta \\ &\quad - \frac{R\tau_c}{4(1+R)^2} (4[(\mathcal{H}^{(\gamma_b)})' + (\mathcal{H}^{(\gamma_b)})^2] v_b \\ &\quad \left. + k[2\mathcal{H}^{(\gamma_b)} \delta_\gamma^{(\gamma_b)} + (\delta_\gamma^{(\gamma_b)})' - 4c_s^2 (\delta_b^{(\gamma_b)})'] \right). \end{aligned} \quad (11)$$

Here the superscript (γ_b) denotes quantities evolved within the modified perturbation system. In particular, $\mathcal{H}$, δ_b , δ_γ , π_γ , and their derivatives are affected indirectly through the modified metric and perturbation evolution. By contrast, the quantities controlling the thermal history of the photon-baryon fluid, namely R , c_s , and τ_c , are left fixed to their background-cosmology values.

With these modifications to the equations directly in the CAMB source code, we obtain the evolution of perturbations as shown in the first panel of Fig. 1. In this panel, all curves correspond to different values of γ_b , while sharing the same cosmological baryon fraction f_b , as defined by the




FIG. 1. Left: redshift evolution of the absolute density contrasts $|\Delta(z)|$ for baryons (dotted lines) and CDM (solid lines) at fixed comoving scale $k = 0.126 \, h \, \text{Mpc}^{-1}$. Three growth-baryon-fraction values are shown: $\gamma_b \approx 0$, $\gamma_b = 0.156$ (fiducial cosmological value), and $\gamma_b = 1$. All lines share identical adiabatic initial conditions and the recombination redshift $z \simeq 1100$. Before decoupling the baryon perturbations undergo acoustic oscillations that are mostly independent of γ_b . After decoupling the growth rate diverges: when $\gamma_b \approx 1$, perturbations evolve solely through baryons, showing delayed growth; when $\gamma_b \approx 0$, perturbations evolve solely through CDM, which grows slightly faster and baryons collapse immediately after recombination. Right: same as on the left, but changing the global baryon fraction f_b as well as the growth-baryon fraction γ_b . The figure shows the scenario where the global baryon fraction is varied for three values: $f_b = \gamma_b \approx 0$, $f_b = \gamma_b = 0.156$ (fiducial), and $f_b = \gamma_b = 1$. While all lines share identical adiabatic initial conditions, they have distinct thermal and growth histories: changing the global baryon fraction not only alters the relative clustering of baryons and CDM, but also affects the recombination epoch and the subsequent evolution of perturbations, leading to significantly different BAO patterns. Since we wish to extract the growth information only while leaving the position of the peaks unchanged, we use the new parameter γ_b to constrain the baryon fraction rather than f_b . Note: the choice of using γ_b or $f_b = 0.011$ as an approximation for the zero baryon fraction case is motivated by the fact that CAMB does not allow a strictly vanishing baryon density. In such a scenario, recombination and reionization cannot be computed, leading to numerical errors. We therefore adopted the smallest allowed value and used it consistently in both plots.

background cosmological parameters. The initial amplitudes of both baryon and CDM perturbations are identical for any value of γ_b , and recombination still occurs at $z \simeq 1100$, as expected given that the growth baryon fraction is not used in computing the initial conditions or the thermal history. The evolution of the perturbation amplitudes, however, changes significantly with γ_b . For baryons, prior to decoupling their oscillatory behavior is set by their tight coupling to photons; because the thermal history is left unchanged with respect to the background cosmology, the baryon sound speed is unaffected, and the predecoupling evolution of baryonic perturbations is therefore essentially independent of γ_b . Instead, after decoupling, they behave quite differently.

To help to visualize this behavior we consider two extreme scenarios.

When $\gamma_b \approx 1$, the evolution of perturbations is driven solely by the baryonic component, enhanced to match the entire matter budget. Since before recombination baryonic perturbations are pressure supported, CDM perturbations do not grow prior to recombination. After decoupling the baryons do not respond to the deep CDM gravitational potential wells present in the fiducial case, and thus we observe in the baryonic perturbations a significant delay

before they begin to grow. Only once they start growing, the CDM perturbations also begin to collapse, reaching the same amplitude as the baryonic perturbations. Although the recombination redshift remains unchanged for all values of γ_b , the residual postrecombination baryonic oscillations—present even in the fiducial case—continue to evolve gravitationally after photon decoupling. In particular, because the gravitational potential is sourced predominantly by the oscillatory baryonic component, acoustic features remains dynamically active for a longer period.

In the opposite scenario, when $\gamma_b \approx 0$, the evolution of perturbations is determined solely by the CDM component. As a result, CDM perturbations grow slightly faster, since they do not respond to the baryonic suppression present before decoupling in the fiducial case. Baryonic perturbations, which remain pressure supported prior to recombination, collapse immediately afterward into the CDM potential wells.

We compare this procedure to that of modifying a single universal baryon fraction in the second panel of Fig. 1. In this panel, the background cosmological parameters are also modified for each curve in order to match f_b to the chosen value of γ_b . We clearly see the effect of the different thermal histories with distinct baryon signature at $z \lesssim 1100$




FIG. 2. Comparison between the linear power spectra obtained with our new method (solid lines) and the KP25 splitting scheme (dashed lines), for different values of γ_b ranging from 0 to 1, including the fiducial cosmological value of 0.156. The two methods encapsulate different physical effects in the definition of the γ_b parameter, leading to visible differences in amplitude, especially at the extremes. At $\gamma_b = 0$, our method removes the baryon-induced suppression on the CDM transfer function prior to the drag epoch. At $\gamma_b = 1$, it also eliminates the CDM-driven growth in baryon oscillations during drag epoch that is retained in the EH transfer functions used by KP25.

leading to differences in the positions of the BAO peaks. Since these modifications include baryonic effects on the thermal history of the Universe and not only on structure growth, they fall outside the scope of this work and are therefore discarded.

Figure 2 shows a comparison between the linear power spectrum obtained with our new approach (solid lines) and that obtained using the previous KP25 splitting scheme (dashed lines), for a range of γ_b values between 0 and 1. For values close to the cosmological baryon fraction, the differences are relatively small, but they become increasingly significant as we move toward either extreme. For $\gamma_b = 0$, the large difference in amplitude comes from the suppression of CDM perturbation growth caused by baryons prior to the drag epoch. When $\gamma_b = 1$, we observe a small difference in the amplitude of the acoustic peaks due to residual CDM-induced growth in the transfer function approach that is not present if we model the different growth at source.

IV. ANALYSIS PIPELINE

Having established a consistent framework to implement the growth baryon fraction γ_b within the perturbation evolution equations, we now turn to its application in the full inference framework of the energy density based H_0 measurement [19]. The application to data from the DESI will be described in our companion paper [30], including the resulting H_0 measurements. In this paper, although we test

the baryon fraction recovery using simulated noiseless theory data vectors only, we follow the procedure used to model the DESI data. In the following, we describe the structure of this analysis, the role of nuisance and cosmological parameters, and the use of HOD-informed priors to mitigate degeneracies and projection effects.

DESI is a redshift survey designed to map the three-dimensional distribution of matter in the Universe with unprecedented precision [36–39]. Operating on the 4-m Mayall Telescope at Kitt Peak National Observatory, DESI uses 5000 robotically controlled optical fibers to obtain spectra of millions of galaxies and quasars across 17,000 deg² of sky [40,41]. Its primary goal is to provide stringent constraints on the expansion history of the Universe and the growth of structure. The first cosmological results, including BAO measurements from the initial year of observations, were presented in 2024 [23,42], and in 2025 the BAO results and cosmological parameter constraints from the second data release were published [43–46].

A. Fitting procedure: Full-shape + compressed postreconstruction BAO

As our baseline analysis, we simultaneously fit the prereconstruction power spectrum multipoles, and the geometric information from the postreconstruction BAO scaling parameters $\alpha_{\parallel}$ and $\alpha_{\perp}$. These parameters quantify the apparent dilation of the BAO feature along and across the line of sight with respect to a fiducial cosmology. They are defined as

$$\alpha_{\parallel} = \frac{D_H(z)/r_d}{[D_H(z)/r_d]_{\text{fid}}}, \quad \alpha_{\perp} = \frac{D_M(z)/r_d}{[D_M(z)/r_d]_{\text{fid}}}, \quad (12)$$

where $D_H(z) \equiv c/H(z)$ is the radial Hubble distance, $D_M(z) \equiv (1+z)D_A(z)$ is the comoving angular diameter distance, and r_d is the sound horizon at the drag epoch. These parameters therefore encode anisotropic geometric distortions in the observed clustering pattern and provide a measurement of the Alcock-Paczynski effect. Since the power spectrum multipoles and the BAO scaling parameters are obtained from the same underlying survey volume, their statistical errors are correlated. We therefore use the joint covariance between the prereconstruction power-spectrum multipoles and the compressed postreconstruction BAO parameters. In practice, we adopt the DESI DR1 covariance structure based on EZmocks, a set of fast approximate mock catalogs designed to reproduce the survey geometry, selection effects, and clustering covariance of the DESI samples [47].

The amplitude of the BAO in the postreconstruction data also depends on γ_b . In our companion paper [30], we tested the impact of incorporating the postreconstruction measurement of γ_b into our pipeline by adding an additional data point and recomputing the covariance accordingly. However, there is no significant improvement in the

TABLE I. Set of priors employed in the baseline analysis for the cosmological parameters and our new density-based method parameters. Uniform and Gaussian priors are denoted by $\mathcal{U}[\text{min}, \text{max}]$ and $\mathcal{N}(\mu, \sigma)$, respectively. The limits for uniform priors correspond to the limits of the training space of the emulator employed in the analysis (for emulator's details see Sec. IV B). The center of the $\omega_{b,\text{dens}}$ prior, μ_{ω_b} , is chosen to be the true value of ω_b for the test under consideration, $\omega_b = 0.02237$ for Abacus and the true ω_b value for the noiseless mock tests.

	Priors
h	$\mathcal{U}[0.1, 1.0]$
$\ln(10^{10} A_s)$	$\mathcal{U}[2.0, 4.0]$
ω_{CDM}	$\mathcal{U}[0.08, 0.16]$
n_s	$\mathcal{U}[0.8, 1.2]$
$\omega_{b,\text{dens}}$	$\mathcal{N}[\mu_{\omega_b}, 0.00055]$
$\Omega_{m,\text{dens}}$	$\mathcal{U}[0.2, 0.5]$
h_{dens}	$\mathcal{U}[0.1, 1.0]$

error on γ_b , since the γ_b constraint from our pipeline and that obtained from postreconstruction BAO for the adopted simulations are correlated at the $\sim 60\%$ level. Consistently, the gain in constraining power is only about $\sim 6\%$. We therefore opted not to include this measurement in our baseline pipeline to avoid introducing additional complexity, particularly regarding the impact of different reconstruction techniques on γ_b .

With the power spectrum multipoles, we constrain the growth baryon fraction γ_b , but also the Λ CDM parameters that shape the broadband component of the linear power spectrum, namely h , $\ln(10^{10} A_s)$, ω_{CDM} , and n_s . The baryon density is only weakly constrained by the power spectrum, so we fix it to $\omega_b = 0.02237$, consistent with the Planck 2018 value [1] adopted in our fiducial cosmology. This choice does not affect our results. The total neutrino mass is fixed to the minimum allowed value $\sum m_\nu = 0.06$ eV,

consistent with the fiducial cosmology. However, in this analysis we are not interested in extracting cosmological information from the broad-band shape, and we marginalize over these other parameters. To fully implement the new density-based method for measuring H_0 , in addition to the growth baryon fraction and the total matter density, we use a measurement of the physical baryon density from BBN. Since these are the measurements employed in our density-based method, and are independent of the values inferred from the broadband information of the power spectrum, we denote them as $\omega_{b,\text{dens}}$ and $\Omega_{m,\text{dens}}$. If we further denote the dimensionless Hubble constant obtained from this method as h_{dens} , we can reparametrize the entire pipeline in terms of these three parameters simply through the following equation

$$\gamma_b = \frac{\omega_{b,\text{dens}}}{\Omega_{m,\text{dens}}(h_{\text{dens}})^2 - \omega_{\text{massive-}\nu}}. \quad (13)$$

As already mentioned, $\Omega_{m,\text{dens}}$ is constrained through BAO dilation parameters, while $\omega_{b,\text{dens}}$ is constrained by imposing a Gaussian BBN prior with a width of 0.00055, matching recent BBN measurements [48]. Finally, we impose flat priors on h_{dens} and $\Omega_{m,\text{dens}}$. In Table I we summarize the priors used for the broadband cosmological parameters and for the new density-based parameters.

The prereconstruction power spectrum multipoles component is modeled with the effective field theory of large-scale structure [49,50], using the implementation in VELOCILEPTORS [51,52]. γ_b modifies only the linear power spectrum, which is then passed to the perturbation-theory calculation. Then the code computes, in both Lagrangian (LPT) and Eulerian (EPT) formalisms, the real- and redshift-space power spectra and correlation functions of biased tracers using 1-loop perturbation theory (including effective field theory counterterms and up to cubic-order biasing). In Fig. 3 we show the effect of γ_b on the monopole




FIG. 3. Comparison of the impact of different values of γ_b on the monopole and quadrupole of the power spectrum computed using EPT for both the KP25 splitting (dashed lines) and our new method (solid lines) using the modified version of CAMB. The different values of γ_b correspond to $\pm 1\sigma$ and $\pm 3\sigma$ around the fiducial value of 0.156.

and quadrupole of the power spectrum, computed with EPT and a fiducial set of EFT parameters, for both our new method and the old KP25 splitting. The variation in γ_b is smaller than in the linear power spectrum and corresponds to $\pm 1\sigma$ and $\pm 3\sigma$ around the fiducial value. We do not show the effect for more extreme values, since the power-spectrum amplitude would enter a regime where the EFT approximation is no longer valid, leading to results that are not reliable and difficult to interpret physically.

The perturbative model introduces a full set of nuisance parameters to account for nonlinear and bias-related effects:

- (i) Bias terms, describing the relation between the galaxy distribution and the underlying matter density field. It includes linear bias b_1 , quadratic bias b_2 , tidal bias b_s , and a third-order bias term b_3 .
- (ii) Counterterms, arising from the effective field theory approach itself, which compensate for the counter effect from the small-scale modes in the perturbative expansion. These are of the form $\alpha_i k^2 P_{\text{lin}}(k)$, with α_i treated as free parameters, usually marginalized over.
- (iii) Stochastic terms, which model shot noise and other contributions such as the Finger-of-God dispersion. These are generally constant or weakly scale-dependent, and may include corrections proportional to k^2 . The amplitude of the stochastic terms is parametrized by the free parameters SN_0 , SN_2 , and SN_4 , which are also marginalized over during the fitting procedure.

An alternative parametrization of the nuisance parameters is also available in VELOCILEPTORS, offering a more physically intuitive approach to prior specification. In the default configuration, the galaxy bias parameters are scaled with $\sigma_8(z)$ as $(1 + b_1)\sigma_8(z)$, $b_2\sigma_8(z)^2$, $b_s\sigma_8(z)^2$, and $b_3\sigma_8(z)^3$. We will refer to this choice as the physical basis, to distinguish it from the standard basis consisting of the unscaled parameters b_1 , b_2 , b_s , and b_3 , i.e. not multiplied by σ_8 . As discussed in [53], this parameter basis more effectively captures the dependence of the monopole on the bias parameters, better aligning with the observables given the high degeneracy between σ_8 and the biases. Other works, such as [54], showed also that this basis provides the best control over prior volume effects when constraining Λ CDM parameters. We ultimately adopted the physical basis as the baseline for our analysis and, since in this work we fit only the monopole and quadrupole of the power spectrum, we fix the nuisance parameters that only affect higher multipoles or are not constrained by this data vector. In particular, we set b_3 , α_4 , α_6 , and SN_4 to zero, consistently with the reduced multipole set used in the analysis. The response of the new parameter γ_b to the choice of bias parametrization is nontrivial. Indeed, in our new approach, the effect of γ_b on the shape of the power spectrum leads to a degeneracy between the new parameter and the EFT nuisances since they are highly degenerate with amplitude

parameters. As the data are already not well constraining the EFT parameters, we can expect even greater projection effects in both the physical and standard bases. To mitigate these effects, we opted for halo occupation distribution (HOD) informed priors on the EFT set of parameters, as presented in [55,56].

Another key aspect relevant for our work in the EFT context is the treatment of infrared (IR) resummation. It is well known that standard 1-loop perturbation theory fails to accurately predict both the position and amplitude of the BAO peak, particularly in real space. This limitation originates from the perturbative expansion itself. Several techniques have therefore been developed to mitigate this effect and properly resum the BAO signal. Our estimate of γ_b is largely driven by the amplitude of the BAO peak, making the impact of IR resummation potentially important for our analysis. We therefore repeated the analysis under different IR-resummation schemes, varying the exponential damping parameter Σ_{BAO}^2 around its fiducial value and modifying the filter used to separate the power spectrum into its wiggled and nonwiggled components. Further details are provided in the companion paper, and we verify that our results are robust to these choices, yielding consistent constraints across all tested configurations.

B. HOD informed priors

To mitigate the strong degeneracies and projection effects associated with poorly constrained nuisance parameters in EFT modeling, we adopt the HOD-informed prior (HIP) approach developed by Zhang *et al.* [56] for DESI. The HIP method uses physically motivated priors on nuisance parameters derived from simulations of galaxy formation based on HOD models. To achieve this, Zhang *et al.* generated real-space mock galaxy catalogs for each DESI tracer using different HOD models within a set of 32 AbacusSummit simulation boxes [57–61], which span a 7-dimensional cosmological parameter space. After converting these catalogs into redshift space and computing the galaxy power spectrum monopole and quadrupole for each realization, they fit the multipoles with the EFT model (specifically, the one-loop Eulerian perturbation theory implementation in VELOCILEPTORS). During the fitting process, the cosmological parameters were kept fixed and so, only the nuisance parameters were fitted. They used these best-fit values to train a normalizing flow model, which learns a highly expressive joint prior distribution over the EFT nuisance parameters. These parameters are expressed in the *physical* basis. These priors were shown to successfully suppress projection effects, particularly in extended cosmological models (such as $w_0 w_a$ CDM), yielding constraints that are consistent with physical expectations and that recover the true cosmology from mock datasets.

However, it is important to note that the simulations used to calibrate these priors were generated under cosmologies




FIG. 4. Validation test of the HOD-informed priors in scenarios where the growth and cosmological baryon fractions differ. The circles show the recovered values of h_{dens} against their true inputs, $h_{\text{dens,true}}$, for both the KP25 splitting (blue) and the new method (red). The black squares show the value of the cosmological Hubble parameter h on the y-axis, as a function of the true h_{dens} value. Their purpose is to illustrate the difference between the h_{dens} value used and the underlying cosmological h , showing that the pipeline successfully recovers the correct h_{dens} with an error smaller than the difference from the cosmological value. The small residual effect of the priors, which slightly push density-based values toward their cosmological counterparts, is negligible compared to the statistical uncertainty (only 10%–15% of the error bar). The shaded gray band highlights the tension in H_0 between Planck 2018 and Pantheon+.

in which the growth baryon fraction matches the cosmological one. As such, it is essential to verify that the application of these priors does not introduce an artificial bias in the inference of γ_b toward the cosmological baryon fraction, especially when this parameter is treated as independent and allowed to vary. To assess this potential issue, we performed an additional dedicated validation test using noiseless data vectors generated from cosmologies in which the growth and cosmological baryon fractions differ (Fig. 4).

The choice to adopt HOD-informed priors comes with a significant performance drawback. Since these priors are not standard distributions but are instead represented as normalizing flows, it is not possible to analytically marginalize over the counterterms and stochastic terms. This requires sampling a fairly high-dimensional parameter space (50 parameters in the case of 6 redshift bins). To optimize the analysis, we therefore implemented the entire pipeline—from theory modeling to likelihood definition and sampling—in JULIA [62], which allowed us to leverage high-performance, state-of-the-art tools. Making use of the same set of best-fit parameters provided by Zhang *et al.*, we retrained the normalizing flows directly in JULIA using the deep learning library LUX.JL [63], and interfaced them with the probabilistic programming language TURING.JL [64,65]

to express our inference model in a compact and efficient way. The use of JULIA also enabled us to take advantage of EFFORT.JL [66], an emulator capable of replicating any PT-based theory code and dramatically accelerating the full analysis. We used it to emulate the EFT model in the Eulerian formulation of VELOCILEPTORS, specifically as implemented in the Resummed Eulerian Perturbation Theory (REPT) class of the official DESI library, DESILIKE.² Additionally, using the ABSTRACTCOSMOLOGICALEMULATORS.JL [66] and EMULATORSTRAINER.JL [66] frameworks, the core of EFFORT.JL emulator, we trained an emulator for $\sigma_8(z)$, since the growth of structures in our model now depends also on the new growth parameters, and an accurate estimate of σ_8 is required to convert EFT parameters from the physical to the standard basis. All these tools support fast and accurate automatic differentiation of every component, allowing us to sample chains with any gradient-based sampler and to obtain best fit solutions using the efficient Limited-memory Broyden–Fletcher–Goldfarb–Shanno algorithm (L-BFGS) minimization algorithm.

V. VALIDATION TESTS

We conducted two main types of validation tests. Having established the presence of projection effects, we first verified that we could adopt the same HOD-informed priors as in [56] without introducing biases in the results. Subsequently, using the final pipeline, we performed a comparison between the previous KP25 splitting scheme and the new method introduced in this work, employing noiseless data vectors generated from a set of Λ CDM (AbacusSummit cosmologies [60,67]) and EDE cosmologies, covering a sufficiently wide range of h values. For both tests, in order to best assess the power of this method at the precision level of currently available data, we employed the DESI DR1 covariance matrices, performing the analysis on the six official DESI redshift bins (three bins for the luminous red galaxies, one for the emission line galaxies, one for the quasars and the bright galaxy sample) [42,47,68,69]. We also restricted the full-shape part of the analysis to the same k range used in DESI Y1, $0.02 \leq k \leq 0.2 \, h \, \text{Mpc}^{-1}$.

We adopted the No-U-Turn Sampler (NUTS) sampler [70] and, to ensure efficient performance and good convergence (Gelman-Rubin criterion of $1 - R < 0.01$), we reparametrized the parameter space and the data vectors such that the variance along each dimension was normalized to unity. This allowed us to run shorter chains while still achieving a convergence of $1 - R < 0.005$ and an average effective sample size (ESS) of at least 1000 for the cosmological parameters, and $1 - R < 0.01$ with an average ESS of at least 800 for the EFT parameters.

²<https://desilike.readthedocs.io/en/latest/>




FIG. 5. Comparison between the true and inferred γ_b and h_{dens} from noiseless data vectors, for both Λ CDM (blue) and EDE (orange) cosmologies. Circles represent the marginalized constraints using the HOD-informed prior, while squares represent the constraints using the default DESI priors. Crosses show the maximum of the posterior. Results are shown for the KP25 splitting method (left) and the new perturbation-evolution approach (right). The bottom panels show the residuals Δ with respect to the true value, with the shaded band marking the Planck–Pantheon+ tension, and the absolute deviation as number of σ (For visual clarity only, we have omitted the error bars for the no HIP case and added a small horizontal shift when necessary).

TABLE II. AbacusSummit cosmologies chosen to test recovery of γ_b .

Cosmo	ω_b	ω_c	h	A_s	n_s	ω_{ncdm}	σ_8
000	0.02237	0.1200	0.6736	2.0830×10^{-9}	0.9649	0.0006442	0.807952
130	0.02237	0.1200	0.6736	1.6140×10^{-9}	0.9649	0.0006442	0.711201
131	0.02237	0.1086	0.7165	2.3146×10^{-9}	0.9649	0.0006442	0.807866
139	0.02416	0.1128	0.7164	2.1681×10^{-9}	0.9732	0.0006442	0.793003
159	0.02206	0.1261	0.5967	1.7977×10^{-9}	0.9673	0.0006442	0.718521
167	0.02248	0.1400	0.7050	1.7321×10^{-9}	0.9715	0.0006442	0.768419
168	0.02157	0.1084	0.6070	2.5378×10^{-9}	0.9275	0.0006442	0.871407
169	0.02189	0.1222	0.6474	1.8067×10^{-9}	0.9898	0.0006442	0.777925
172	0.02282	0.1032	0.6369	2.1259×10^{-9}	0.9012	0.0006442	0.765650
177	0.02239	0.1344	0.6820	2.0641×10^{-9}	1.0002	0.0006442	0.806026
181	0.02169	0.1112	0.5745	2.0651×10^{-9}	0.9336	0.0006442	0.730036

A. HIP validation

We verified the validity of the HOD priors in scenarios where the growth and cosmological baryon fractions do not match. We selected a set of simulations generated from different cosmologies in which the cosmological and growth baryon fractions coincide. We then performed a fit of all parameters except for γ_b , which was fixed to a value different from the cosmological baryon fraction. We fixed γ_b to values up to 27% different from the cosmological baryon fraction, which translates into h_{dens} values up to 16% different from the cosmological h . The resulting fit provides a set of cosmological and EFT parameters giving approximately the same power in the multipoles. Using these parameters, we generated new noiseless simulations of a nonphysical Universe in which the growth and cosmological baryon fractions no longer coincide. This is equivalent to fixing σ_8 to a different value and adjusting the other parameters accordingly. As shown in Fig. 4, the pipeline is able to recover the true h_{dens} used to generate the data, even when this value is, in some cases, significantly different from the cosmological one.

It is also evident that there exists a correlation between the bias and the cosmological Hubble parameter: when the cosmological Hubble parameter h exceeds h_{dens} , the resulting bias is positive; conversely, when it is lower, the bias becomes negative. This convinced us that the HIPs have a minor effect on the γ_b recovered from our pipeline, slightly pushing the values toward their cosmological counterparts. This effect is very small and, relative to the error bars, is negligible, amounting to only about 10%–15% of the standard deviation. For clarity, we also overplot a shaded gray area representing the tension between the H_0 values from Planck 2018 [1] and Pantheon+ [71]. Similarly, we verified that both γ_b and $\Omega_{m,\text{dens}}$ are unbiased, obtaining the same correlation between the sign of the residual bias and the cosmological parameter, but again with the bias that remains within the same 10%–15% of the corresponding σ .

This confirms that the information encoded in the HOD priors does not induce spurious constraints or biases on

these parameters, and can therefore be safely employed within our energy density framework.

B. Noiseless data vectors test

We have conducted a series of validations on noiseless data vectors generated from a set of Λ CDM and EDE cosmologies. In this analysis, we applied both the previous KP25 splitting scheme and the new method introduced in this work, in order to carefully assess their differences in terms of the precision with which they recover the value of the Hubble constant, and the residual bias from projection effects. We selected the Λ CDM cosmologies from the AbacusSummit suite, specifically the 10 listed in Table II. For the EDE case, we fixed the fiducial cosmological parameters as studied in [7] ($h = 0.7219$, $\omega_b = 0.02253$, $\omega_c = 0.1306$, $A_s = 2.215 \times 10^{-9}$, $n_s = 0.9889$, $f_{\text{EDE}} = 0.122$, $\log_{10} z_c = 3.562$, $\theta_{i,\text{scf}} = 2.83$), and then considered nine variations of this cosmology by varying h while keeping ω_b and Ω_m fixed, so that changes in h would fully translate into variations of the baryon fraction. With the Λ CDM cosmologies we covered the h range 0.57–0.72, while the EDE cosmologies spanned the range 0.66–0.77. In Fig. 5, we compare the true value of γ_b and h with the value inferred by our pipeline for the noiseless Λ CDM and EDE data vectors. We find that the two methods yield equally precise results, corresponding to relative marginalized 1σ uncertainties of approximately $\sim 8\%$ for γ_b and $\sim 5\%$ for h_{dens} . However, the KP25 method appears to retain a slightly larger bias compared to the new approach. The resulting Δh and $\Delta \gamma_b$ values remain entirely negligible compared to the constraining power of current data for both methods, as well as to the most relevant H_0 tension. We additionally plot the biases in γ_b and h in units of their error. When using the default prior on the biases and counterterms, we find significant projection effects leading to $\sim 0.5\sigma$ biases in the marginalized posteriors. The HOD-informed prior effectively mitigates these projection effects, reducing the biases on the marginalized means back to the $\lesssim 0.1\sigma$ range. Thus, we have proven that the new

approach is capable of inferring the correct value of γ_b , and therefore H_0 , regardless of whether the Universe is Λ CDM or EDE.

The old splitting and our new method both extract their primary information from the BAO amplitude, but they differ in how the baryon fraction affects the smooth shape of the power spectrum. The two approaches are therefore complementary, and the fact that they yield consistent results at this level of precision provides a confirmation of the robustness of this approach.

VI. CONCLUSIONS

In this paper we have developed a new framework to measure the baryon fraction from galaxy clustering, with the explicit goal of enabling energy density based determinations of the Hubble constant. Previous work introduced the idea of extracting the baryon fraction from the amplitude of the baryon signal in the galaxy power spectrum, but implemented it through an approximate template-based transfer-function splitting that neglected part of the baryon contribution to the growth of structure.

Here, we have proposed an improved, physically self-consistent method that embeds the growth baryon fraction parameter γ_b directly into the linear perturbation evolution equations of CAMB. This ensures that both baryons and CDM respond coherently to variations in γ_b across all cosmic epochs, while preserving the correct thermal history. We integrated this implementation into a complete EFT-based full-shape analysis pipeline, further enhanced with HOD-informed priors to mitigate degeneracies and projection effects.

Through validation tests on noiseless data vectors from both Λ CDM and EDE cosmologies, we demonstrated that the new approach achieves comparable precision to the previous splitting scheme (about 8% on γ_b and 5% on h), while substantially reducing systematic biases. Importantly, it maintains the ability to recover the correct value of γ_b and thus H_0 even in extended cosmologies, confirming its robustness. Although the statistical precision is not improved relative to the previous method, the new approach offers a cleaner and more accurate physical description and therefore represents a more reliable foundation for future applications. Overall, our results show that incorporating γ_b at the perturbation level provides a robust and self-consistent route to baryon-fraction measurements. Combined with BBN constraints on ω_b and Alcock-Paczynski determinations of Ω_m , this establishes a viable way to measure H_0 independently of the sound horizon.

The application of this method already represents a significant innovation in the context of the H_0 tension, with substantial prospects for improvement. Since the precision on H_0 is entirely determined by the precision on γ_b , which improves with increasing survey volume, this methodology will have the opportunity to demonstrate its strength in ongoing and future deep surveys such as DESI,

Euclid [72,73], Rubin Observatory's Legacy Survey of Space and Time (LSST) [74], and Roman [75]. DESI will cover a comoving volume of $\sim 60 h^{-3} \text{ Gpc}^3$ [76], nearly ten times that of BOSS, reducing the statistical uncertainty on γ_b significantly and enabling H_0 determinations precise enough to clearly separate the SH0ES and CMB/LSS values at $> 3\sigma$. Comparable sensitivity is expected from Euclid, with a volume of $\sim 55 h^{-3} \text{ Gpc}^3$ [73], while additional progress will come from all-sky spectroscopic surveys such as SPHEREx [77] and the Roman Space Telescope [78]. Meanwhile, BAO or full-shape-based analyses tied to r_d are forecast to reach subpercent precision, with uncertainties below $\sigma_{H_0} < 0.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [15,79]. The combination of these complementary methods will provide stringent cross-checks, offering a decisive opportunity to test the robustness of Λ CDM and to explore possible new physics behind the H_0 discrepancy.

ACKNOWLEDGMENTS

We thank Antony Lewis for useful conversations about the structure of CAMB. A. K. was supported as a CITA National Fellow by the Natural Sciences and Engineering Research Council of Canada (NSERC), funding reference No. DIS-2022-568580. W. P. acknowledges support from the Natural Sciences and Engineering Research Council of Canada (NSERC), [funding reference No. RGPIN-2025-03931] and from the Canadian Space Agency. Research at Perimeter Institute is supported in part by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Colleges and Universities. This research was enabled in part by support provided by Compute Ontario (computeontario.ca) and the Digital Research Alliance of Canada (alliancecan.ca). This material is based upon work supported by the U.S. Department of Energy (DOE), Office of Science, Office of High-Energy Physics, under Contract No. DE-AC02-05CH11231, and by the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility under the same contract. Additional support for DESI was provided by the U.S. National Science Foundation (NSF), Division of Astronomical Sciences under Contract No. AST-0950945 to the NSF's National Optical-Infrared Astronomy Research Laboratory; the Science and Technology Facilities Council of the United Kingdom; the Gordon and Betty Moore Foundation; the Heising-Simons Foundation; the French Alternative Energies and Atomic Energy Commission (CEA); the National Council of Humanities, Science and Technology of Mexico (CONAHCYT); the Ministry of Science, Innovation and Universities of Spain (MICIU/AEI/10.13039/501100011033), and by the DESI Member Institutions: [80]. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the

U.S. National Science Foundation, the U.S. Department of Energy, or any of the listed funding agencies. The authors are honored to be permitted to conduct scientific research on I'oligam Du'ag (Kitt Peak), a mountain with particular significance to the Tohono O'odham Nation.

DATA AVAILABILITY

The data used in this work are public as part of DESI Data Release 1 [81]. The data points corresponding to the figures are available on Zenodo [82]. Code is available on github [83].

-
- [1] N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi *et al.* (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, *Astron. Astrophys.* **641**, A2020 (2020).
 - [2] A. G. Riess, W. Yuan, L. M. Macri, D. Scolnic, D. Brout, S. Casertano *et al.*, A comprehensive measurement of the local value of the Hubble constant with $1 \text{ km s}^{-1} \text{ Mpc}^{-1}$ uncertainty from the Hubble space telescope and the SH0ES Team, *Astrophys. J. Lett.* **934**, L2022 (2022).
 - [3] R. J. Cooke, M. Pettini, K. M. Nollett, and R. Jorgenson, The primordial deuterium abundance of the most metal-poor damped Lyman- α system, *Astrophys. J.* **830**, 148 (2016).
 - [4] T. Karwal and M. Kamionkowski, Dark energy at early times, the Hubble parameter, and the string axiverse, *Phys. Rev. D* **94**, 103523 (2016).
 - [5] V. Poulin, T. L. Smith, T. Karwal, and M. Kamionkowski, Early dark energy can resolve the Hubble tension, *Phys. Rev. Lett.* **122**, 221301 (2019).
 - [6] T. L. Smith, V. Poulin, and M. A. Amin, Oscillating scalar fields and the Hubble tension: A resolution with novel signatures, *Phys. Rev. D* **101**, 063523 (2020).
 - [7] J. C. Hill, E. McDonough, M. W. Toomey, and S. Alexander, Early dark energy does not restore cosmological concordance, *Phys. Rev. D* **102**, 043507 (2020).
 - [8] G. P. Lynch, L. Knox, and J. Chluba, DESI observations and the Hubble tension in light of modified recombination, *Phys. Rev. D* **110**, 083538 (2024).
 - [9] S. H. Mirpoorian, K. Jedamzik, and L. Pogosian, Modified recombination and the Hubble tension, *Phys. Rev. D* **111**, 083519 (2025).
 - [10] N. Schöneberg and L. Vacher, The mass effect—variations of the electron mass and their impact on cosmology, *J. Cosmol. Astropart. Phys.* **03** (2025) 004.
 - [11] Atacama Cosmology Telescope Collaboration, The Atacama Cosmology Telescope: DR6 constraints on extended cosmological models, *J. Cosmol. Astropart. Phys.* **11** (2025) 063.
 - [12] V. Poulin, T. L. Smith, R. Calderón, and T. Simon, Impact of ACT DR6 and DESI DR2 for early dark energy and the Hubble tension, [arXiv:2505.08051](https://arxiv.org/abs/2505.08051).
 - [13] O. H. E. Philcox, G. S. Farren, B. D. Sherwin, E. J. Baxter, and D. J. Brout, Determining the Hubble constant without the sound horizon: A 3.6% constraint on H_0 from galaxy surveys, CMB lensing, and supernovae, *Phys. Rev. D* **106**, 063530 (2022).
 - [14] E. J. Baxter and B. D. Sherwin, Determining the Hubble constant without the sound horizon scale: Measurements from CMB lensing, *Mon. Not. R. Astron. Soc.* **501**, 1823 (2021).
 - [15] G. S. Farren, O. H. E. Philcox, and B. D. Sherwin, Determining the Hubble constant without the sound horizon: Perspectives with future galaxy surveys, *Phys. Rev. D* **105**, 063503 (2022).
 - [16] O. H. E. Philcox, B. D. Sherwin, G. S. Farren, and E. J. Baxter, Determining the Hubble constant without the sound horizon: Measurements from galaxy surveys, *Phys. Rev. D* **103**, 023538 (2021).
 - [17] B. Bahr-Kalus, D. Parkinson, K. Lodha, E. Mueller, E. Chaussidon, A. de Mattia *et al.*, Model-independent measurement of the matter-radiation equality scale in DESI 2024, *Phys. Rev. D* **112**, 063553 (2025).
 - [18] E. Zaborowski, P. Taylor, K. Honscheid, A. Cuceu, A. de Mattia, D. Huterer *et al.*, A sound horizon-free measurement of H_0 in DESI 2024, *J. Cosmol. Astropart. Phys.* **06** (2025) 020.
 - [19] A. Krolewski, W. J. Percival, and A. Woodfinden, New method to determine the Hubble parameter from cosmological energy-density measurements, *Phys. Rev. Lett.* **134**, 101002 (2025).
 - [20] G. Lavaux and B. D. Wandelt, Precision cosmography with stacked voids, *Astrophys. J.* **754**, 109 (2012).
 - [21] A. Krolewski and W. J. Percival, Measuring the baryon fraction using galaxy clustering, *Phys. Rev. D* **111**, 063526 (2025).
 - [22] DESI Collaboration, The DESI experiment part I: Science, targeting, and survey design, [arXiv:1611.00036](https://arxiv.org/abs/1611.00036).
 - [23] D. Collaboration, M. Abdul-Karim, A. G. Adame, D. Aguado, J. Aguilar, S. Ahlen *et al.*, Data release 1 of the dark energy spectroscopic instrument, [arXiv:2503.14745](https://arxiv.org/abs/2503.14745).
 - [24] S. Alam, M. Aubert, S. Avila, C. Bolland, J. E. Bautista, M. A. Bershad *et al.*, Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory, *Phys. Rev. D* **103**, 083533 (2021).
 - [25] D. J. Eisenstein and W. Hu, Baryonic features in the matter transfer function, *Astrophys. J.* **496**, 605 (1998).
 - [26] J. Lesgourgues, The cosmic linear anisotropy solving system (class) I: Overview, [arXiv:1104.2932](https://arxiv.org/abs/1104.2932).
 - [27] C. Heinrich, Lensing bias to CMB polarization measurements of compensated isocurvature perturbations, *Phys. Rev. D* **97**, 023513 (2018).
 - [28] A. Barreira, G. Cabass, K. D. Lozanov, and F. Schmidt, Compensated isocurvature perturbations in the galaxy power spectrum, *J. Cosmol. Astropart. Phys.* **07** (2020) 049.

- [29] A. Barreira, Constraints on compensated isocurvature perturbations from BOSS DR12 galaxy data, [arXiv:2302.01927](#).
- [30] A. Krolewski, A. Crespi, and W. Percival, A measurement of H_0 from DESI DR1 using energy densities, [arXiv:2511.23432](#).
- [31] A. Lewis, A. Challinor, and A. Lasenby, Efficient computation of cosmic microwave background anisotropies in closed Friedmann-Robertson-Walker models, *Astrophys. J.* **538**, 473 (2000).
- [32] U. Andrade, D. Anbajagane, R. von Martens, D. Huterer, and J. Alcaniz, A test of the standard cosmological model with geometry and growth, *J. Cosmol. Astropart. Phys.* **11** (2021) 014.
- [33] M. Chu and L. Knox, Testing cosmological models and understanding cosmological parameter determinations with metaparameters, *Astrophys. J.* **620**, 1 (2005).
- [34] A. M. Lewis, Geometric algebra and covariant methods in physics and cosmology, Ph.D. thesis, Apollo—University of Cambridge Repository, [10.17863/CAM.16632](#) (2001).
- [35] A. Lewis, CAMB Notes, <https://cosmologist.info/notes/CAMB.pdf>.
- [36] A. Aghamousa, J. Aguilar, S. Ahlen, S. Alam, L. E. Allen, C. A. Prieto *et al.*, The DESI experiment part II: Instrument design, [arXiv:1611.00037](#).
- [37] B. Abareschi, J. Aguilar, S. Ahlen, S. Alam, D. M. Alexander, R. Alfarsy *et al.*, Overview of the instrumentation for the dark energy spectroscopic instrument, *Astron. J.* **164**, 207 (2022).
- [38] T. N. Miller, P. Doel, G. Gutierrez, R. Besuner, D. Brooks, G. Gallo *et al.*, The optical corrector for the dark energy spectroscopic instrument, *Astron. J.* **168**, 95 (2024).
- [39] C. Poppett, L. Tyas, J. Aguilar, C. Bebek, D. Bramall, T. Claybaugh *et al.*, Overview of the fiber system for the dark energy spectroscopic instrument, *Astron. J.* **168**, 245 (2024).
- [40] J. Guy, S. Bailey, A. Kremin, S. Alam, D. M. Alexander, C. Allende Prieto *et al.*, The spectroscopic data processing pipeline for the dark energy spectroscopic instrument, *Astron. J.* **165**, 144 (2023).
- [41] E. F. Schlafly, D. Kirkby, D. J. Schlegel, A. D. Myers, A. Raichoor, K. Dawson *et al.*, Survey operations for the dark energy spectroscopic instrument, *Astron. J.* **166**, 259 (2023).
- [42] A. Adame, J. Aguilar, S. Ahlen, S. Alam, D. Alexander, C. Allende Prieto *et al.*, DESI 2024 VII: Cosmological constraints from the full-shape modeling of clustering measurements, *J. Cosmol. Astropart. Phys.* **07** (2025) 028.
- [43] M. Abdul-Karim, J. Aguilar, S. Ahlen, S. Alam, L. Allen, C. A. Prieto *et al.*, DESI DR2 results. II. Measurements of baryon acoustic oscillations and cosmological constraints, *Phys. Rev. D* **112**, 083515 (2025).
- [44] DESI Collaboration, DESI DR2 results. I. Baryon acoustic oscillations from the Lyman alpha forest, *Phys. Rev. D* **112**, 083514 (2025).
- [45] DESI Collaboration, Validation of the DESI DR2 measurements of baryon acoustic oscillations from galaxies and quasars, *Phys. Rev. D* **112**, 083512 (2025).
- [46] L. Casas, H. K. Herrera-Alcantar, J. Chaves-Montero, A. Cuceu, A. Font-Ribera, M. Lokken *et al.*, Validation of the DESI DR2 Ly α BAO analysis using synthetic datasets, [arXiv:2503.14741](#).
- [47] D. Forero-Sánchez, M. Rashkovetskyi, O. Alves, A. de Mattia, N. Padmanabhan, H. Seo *et al.*, Analytical and EZMOCK covariance validation for the DESI 2024 results, *J. Cosmol. Astropart. Phys.* **04** (2025) 055.
- [48] N. Schöneberg, The 2024 BBN baryon abundance update, *J. Cosmol. Astropart. Phys.* **06** (2024) 006.
- [49] D. Baumann, A. Nicolis, L. Senatore, and M. Zaldarriaga, Cosmological nonlinearities as an effective fluid, *J. Cosmol. Astropart. Phys.* **07** (2012) 051.
- [50] J. J. M. Carrasco, M. P. Hertzberg, and L. Senatore, The effective field theory of cosmological large scale structures, *J. High Energy Phys.* **09** (2012) 082.
- [51] S.-F. Chen, Z. Vlah, and M. White, Consistent modeling of velocity statistics and redshift-space distortions in one-loop perturbation theory, *J. Cosmol. Astropart. Phys.* **07** (2020) 062.
- [52] S.-F. Chen, Z. Vlah, E. Castorina, and M. White, Redshift-space distortions in Lagrangian perturbation theory, *J. Cosmol. Astropart. Phys.* **03** (2021) 100.
- [53] M. Maus, S. Chen, M. White, J. Aguilar, S. Ahlen, A. Aviles *et al.*, An analysis of parameter compression and Full-Modeling techniques with Velocileptors for DESI 2024 and beyond, *J. Cosmol. Astropart. Phys.* **01** (2025) 138.
- [54] M. Maus, Y. Lai, H. E. Noriega, S. Ramirez-Solano, A. Aviles, S. Chen *et al.*, A comparison of effective field theory models of redshift space galaxy power spectra for DESI 2024 and future surveys, *J. Cosmol. Astropart. Phys.* **01** (2025) 134.
- [55] H. Zhang, M. Bonici, G. D’Amico, S. Paradiso, and W. J. Percival, HOD-informed prior for EFT-based full-shape analyses of LSS, *J. Cosmol. Astropart. Phys.* **04** (2025) 041.
- [56] DESI Collaboration, Enhancing DESI DR1 full-shape analyses using HOD-informed priors, *J. Cosmol. Astropart. Phys.* **11** (2025) 049.
- [57] L. H. Garrison, D. J. Eisenstein, D. Ferrer, M. V. Metchnik, and P. A. Pinto, Improving initial conditions for cosmological n -body simulations, *Mon. Not. R. Astron. Soc.* **461**, 4125 (2016).
- [58] L. H. Garrison, D. J. Eisenstein, D. Ferrer, J. L. Tinker, P. A. Pinto, and D. H. Weinberg, The abacus cosmos: A suite of cosmological N -body simulations, *Astrophys. J. Suppl. Ser.* **236**, 43 (2018).
- [59] L. H. Garrison, D. J. Eisenstein, and P. A. Pinto, A high-fidelity realization of the euclid code comparison N -body simulation with abacus, *Mon. Not. R. Astron. Soc.* **485**, 3370 (2019).
- [60] L. H. Garrison, D. J. Eisenstein, D. Ferrer, N. A. Maksimova, and P. A. Pinto, The abacus cosmological N -body code, *Mon. Not. R. Astron. Soc.* **508**, 575 (2021).
- [61] S. Bose, D. J. Eisenstein, B. Hadzhiyska, L. H. Garrison, and S. Yuan, Constructing high-fidelity halo merger trees in abacussummit, *Mon. Not. R. Astron. Soc.* **512**, 837 (2022).
- [62] J. Bezanson, A. Edelman, S. Karpinski, and V. B. Shah, Julia: A fresh approach to numerical computing, [arXiv:1411.1607](#).
- [63] A. Pal, Lux: Explicit parameterization of deep neural networks in Julia, [10.5281/zenodo.7808903](#) (2023).
- [64] T. E. Fjelde, K. Xu, D. Widmann, M. Tarek, C. Pfiffer, M. Trapp *et al.*, Turing.jl: A general-purpose probabilistic

- programming language, *ACM Trans. Probab. Mach. Learn.* **1**, 14 (2025).
- [65] H. Ge, K. Xu, and Z. Ghahramani, Turing: A language for flexible probabilistic inference, in *Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics*, edited by A. Storkey and F. Perez-Cruz, vol. 84 of *Proceedings of Machine Learning Research* (PMLR, 2018), pp. 1682–1690, <https://proceedings.mlr.press/v84/ge18b.html>.
- [66] M. Bonici, G. D’Amico, J. Bel, and C. Carbone, Effort.jl: A fast and differentiable emulator for the effective field theory of the large scale structure of the universe, *J. Cosmol. Astropart. Phys.* **09** (2025) 044.
- [67] N. A. Maksimova, L. H. Garrison, D. J. Eisenstein, B. Hadzhiyska, S. Bose, and T. P. Satterthwaite, AbacusSummit: A massive set of high-accuracy, high-resolution N-body simulations, *Mon. Not. R. Astron. Soc.* **508**, 4017 (2021).
- [68] DESI Collaboration, DESI 2024 V: Full-Shape galaxy clustering from galaxies and quasars, *J. Cosmol. Astropart. Phys.* **09** (2025) 008.
- [69] M. Rashkovetskyi, D. Forero-Sanchez, A. de Mattia, D. Eisenstein, N. Padmanabhan, H. Seo *et al.*, Semianalytical covariance matrices for two-point correlation function for DESI 2024 data, *J. Cosmol. Astropart. Phys.* **01** (2025) 145.
- [70] M. D. Hoffman and A. Gelman, The no-u-turn sampler: Adaptively setting path lengths in Hamiltonian Monte Carlo, [arXiv:1111.4246](https://arxiv.org/abs/1111.4246).
- [71] D. Brout, D. Scolnic, B. Popovic, A. G. Riess, A. Carr, J. Zuntz *et al.*, The pantheon+analysis: Cosmological constraints, *Astrophys. J.* **938**, 110 (2022).
- [72] Y. Mellier, Abdurro’uf, J. A. Acevedo Barroso, A. Achucarro, J. Adamek, R. Adam *et al.*, Euclid: I. Overview of the euclid mission, *Astron. Astrophys.* **697**, A2025 (2025).
- [73] A. Blanchard, S. Camera, C. Carbone, V. F. Cardone, S. Casas *et al.* (Euclid Collaboration), Euclid preparation. VII. Forecast validation for Euclid cosmological probes, *Astron. Astrophys.* **642**, A2020 (2020).
- [74] v. Ivezić, S. M. Kahn, J. A. Tyson, B. Abel, E. Acosta, R. Allsman *et al.*, Lsst: From science drivers to reference design and anticipated data products, *Astrophys. J.* **873**, 111 (2019).
- [75] G. Mosby, B. J. Rauscher, C. Bennett, E. S. Cheng, S. Cheung, A. Cillis *et al.*, Properties and characteristics of the Nancy Grace Roman Space Telescope H4RG-10 detectors, *J. Astron. Telesc. Instrum. Syst.* **6**, 046001 (2020).
- [76] A. G. Adame, J. Aguilar, S. Ahlen, S. Alam, G. Aldering, D. M. Alexander *et al.*, Validation of the scientific program for the dark energy spectroscopic instrument, *Astron. J.* **167**, 62 (2024).
- [77] O. Doré, J. Bock, M. Ashby, P. Capak, A. Cooray, R. de Putter *et al.*, Cosmology with the SPHEREX all-sky spectral survey, [arXiv:1412.4872](https://arxiv.org/abs/1412.4872).
- [78] T. Eifler, H. Miyatake, E. Krause, C. Heinrich, V. Miranda, C. Hirata *et al.*, Cosmology with the Roman space telescope—multiprobe strategies, *Mon. Not. R. Astron. Soc.* **507**, 1746 (2021).
- [79] M. M. Ivanov and O. H. E. Philcox, Measuring H_0 with spectroscopic surveys, [arXiv:2305.07977](https://arxiv.org/abs/2305.07977).
- [80] <https://www.desi.lbl.gov/collaborating-institutions>
- [81] DESI Collaboration, DESI Data Release 1, <https://data.desi.lbl.gov/doc/releases/> (2025).
- [82] A. Krolewski, A measurement of H_0 from DESI DR1 using energy densities, [10.5281/zenodo.17686137](https://doi.org/10.5281/zenodo.17686137) (2025).
- [83] A. Crespi, fbCAMB: Modified CAMB with a growth baryon-fraction parameter, <https://github.com/drew2799/fbCAMB> (2026).