

Lecture 4

Superspace

Outline

- Geometric realization of supersymmetry: superspace
- General scalar superfields.
- Examples: chiral superfield, vector superfield, the (spinorial) field strength superfield.
- Extended gauge invariance, and the WZ-gauge.
- Lagrangians in superspace.

Reading: Terning 2.7, Lykken's TASI notes (arXiv: 9612114), sec 3.

Superspace

The idea: generalize the geometric interpretation of the Lorentz group, as the symmetry of 4D Minkowski space.

Implementation: supersymmetry is the symmetry of **superspace**, an extension of 4D Minkowski space to include additional “fermionic” directions.

Superspace coordinates: $(x^\mu, \theta_\alpha, \bar{\theta}_{\dot{\alpha}})$ where $\theta_\alpha, \bar{\theta}_{\dot{\alpha}} = \theta_\alpha^\dagger$ are **anticommuting** (Grassmann) two-component spinors satisfying

$$\{\theta_\alpha, \bar{\theta}_{\dot{\alpha}}\} = 0$$

Supersymmetry is the symmetry group of $(x^\mu, \theta_\alpha, \bar{\theta}_{\dot{\alpha}})$. The Lorentz subgroup acts according to the Lorentz indices, but other SUSY generators mix bosonic and fermionic coordinates.

Integrals over Superspace

Reminder: for a single component Grassmann variable η , integrals are defined so:

$$\int d\eta = 0, \quad \int d\eta \, \eta = 1$$

For chiral superspace coordinates define:

$$d^2\theta \equiv -\frac{1}{4}d\theta^\alpha d\theta^\beta \epsilon_{\alpha\beta}$$

Then

$$\begin{aligned} \int d^2\theta \, \theta^2 &= -\frac{1}{4} \int d\theta^\alpha d\theta^\beta \epsilon_{\alpha\beta} (\theta^\sigma \epsilon_{\sigma\tau} \theta^\tau) \\ &= -\frac{1}{4} \epsilon_{\alpha\beta} \epsilon_{\sigma\tau} (\delta^{\beta\sigma} \delta^{\tau\alpha} - \delta^{\alpha\sigma} \delta^{\tau\beta}) \\ &= -\frac{1}{2} \epsilon_{\alpha\beta} \epsilon_{\beta\alpha} = 1 \end{aligned}$$

Similarly, for arbitrary spinors χ and ψ :

$$\int d^2\theta (\chi\theta)(\psi\theta) = -\frac{1}{2}(\chi\psi)$$

The analogous superspace measure for the conjugate superspace coordinate:

$$d^2\bar{\theta} \equiv -\frac{1}{4}d\bar{\theta}_{\dot{\alpha}}d\bar{\theta}_{\dot{\beta}}\epsilon^{\dot{\alpha}\dot{\beta}}$$

Notation for the nonchiral measure over all of the fermionic coordinates

$$d^4\theta \equiv d^2\theta d^2\bar{\theta}$$

The measure over all of superspace

$$d^4x d^4\theta$$

It is this measure that is invariant under all SUSY transformations.

General Scalar Superfield

Component expansion of scalar field on superspace (**warning**: notation does not not agree in detail with ours):

$$\begin{aligned}\Phi(x, \theta, \bar{\theta}) = & f(x) + \theta\phi(x) + \bar{\theta}\bar{\chi}(x) + \theta^2 m(x) + \bar{\theta}^2 n(x) \\ & + \theta\sigma^\mu\bar{\theta}v_\mu(x) + \theta^2\bar{\theta}\bar{\lambda}(x) + \bar{\theta}^2\theta\psi(x) + \theta^2\bar{\theta}^2 d(x)\end{aligned}$$

The component fields $f(x)$, $m(x)$, $n(x)$, $v_\mu(x)$, $d(x)$ are complex bosonic fields.

The component fields $\phi(x)$, $\bar{\chi}(x)$, $\bar{\lambda}(x)$, $\psi(x)$ are fermionic fields that anti-commute with each other and the superspace coordinates $\theta, \bar{\theta}$.

There are $2+2+2+8+2 = 16$ bosonic components and $4+4+4+4 = 16$ fermionic components.

Superspace transformations

Goal: realize the superalgebra on superspace.

Translate (left-multiply) by $G(y, \epsilon, \bar{\epsilon})$ on the coset element $\Omega(x, \theta, \bar{\theta})$ (=“the wave function”):

$$\begin{aligned} G(y, \epsilon, \bar{\epsilon})\Omega(x, \theta, \bar{\theta}) &= e^{i(-y^\mu P_\mu + \epsilon Q + \bar{\epsilon}\bar{Q})} e^{i(-x^\mu P_\mu + \theta Q + \bar{\theta}\bar{Q})} \\ &= e^{i(-(x^\mu + y^\mu)P_\mu + (\theta + \epsilon)Q + (\bar{\epsilon} + \bar{\theta})\bar{Q} + \frac{i}{2}[\epsilon Q, \bar{\theta}\bar{Q}] + \frac{i}{2}[\bar{\epsilon}\bar{Q}, \theta Q])} \\ &= \Omega(x^\mu + y^\mu - i\epsilon\sigma^\mu\bar{\theta} + i\theta\sigma^\mu\bar{\epsilon}, \theta + \epsilon, \bar{\theta} + \bar{\epsilon}) \end{aligned}$$

Intermediate steps

- The Baker-Campbell-Hausdorff formula (the ...-terms vanish presently):

$$e^A e^B = e^{A+B+\frac{1}{2}[A,B]+\dots}$$

- The fundamental anticommutator in the form

$$[\epsilon Q, \bar{\theta}\bar{Q}] = 2\epsilon\sigma^\mu\bar{\theta}P_\mu$$

The result: the translation generator and the supertranslation generators are realized on superspace as

$$\begin{aligned} i\partial_\mu &= P_\mu \\ Q_\alpha &= -i \left(\partial_\alpha + i\sigma_{\alpha\dot{\beta}}^\mu \bar{\theta}^{\dot{\beta}} \partial_\mu \right) \\ Q_{\dot{\alpha}}^\dagger &= i \left(\bar{\partial}_{\dot{\alpha}} + i\theta^\beta \sigma_{\beta\dot{\alpha}}^\mu \partial_\mu \right) \end{aligned}$$

where (as usual) the spinorial derivatives:

$$\begin{aligned} \partial_\alpha &= \frac{\partial}{\partial \theta^\alpha} \\ \bar{\partial}_{\dot{\alpha}} &= \frac{\partial}{\partial \bar{\theta}^{\dot{\alpha}}} \end{aligned}$$

act **on the left**.

Examples:

$$\begin{aligned} \partial_\alpha \theta^2 &= \partial_\alpha (\theta^\beta \theta_\beta) = 2\theta_\alpha \\ \bar{\partial}_{\dot{\alpha}} \bar{\theta}^2 &= \bar{\partial}_{\dot{\alpha}} (\bar{\theta}_{\dot{\beta}} \bar{\theta}^{\dot{\beta}}) = -2\theta_{\dot{\alpha}} \end{aligned}$$

SUSY transformation of superfield

The SUSY transform of a general scalar superfield:

$$\begin{aligned}\delta_\epsilon \Phi(x, \theta, \bar{\theta}) &= i(\epsilon Q + \epsilon^\dagger Q^\dagger) \Phi(x, \theta, \bar{\theta}) \\ &= \epsilon \phi + \bar{\epsilon} \bar{\chi} + i\theta \sigma^\mu \bar{\epsilon} \partial_\mu f + 2\epsilon \theta m + \theta \sigma^\mu \bar{\epsilon} v_\mu - i\epsilon \sigma^\mu \bar{\theta} \partial_\mu f \\ &+ 2\bar{\epsilon} \bar{\theta} n + \epsilon \sigma^\mu \bar{\theta} v_\mu + i(\theta \sigma^\mu \epsilon) \theta \partial_\mu \phi + \theta^2 \bar{\epsilon} \bar{\lambda} - i(\epsilon \sigma^\mu \bar{\theta}) \bar{\theta} \sigma_\mu \bar{\chi} \\ &+ \bar{\theta}^2 \epsilon \psi - i\epsilon \sigma^\mu \bar{\theta} \theta \partial_\mu \phi + i\theta \sigma^\mu \bar{\epsilon} \bar{\theta} \partial_\mu \bar{\chi} + 2\epsilon \theta \bar{\theta} \bar{\lambda} + 2\bar{\epsilon} \bar{\theta} \theta \psi \\ &- i\epsilon \sigma^\mu \bar{\theta} \theta^2 \partial_\mu m + i\theta \sigma^\mu \bar{\epsilon} \theta \sigma^\nu \bar{\theta} \partial_\mu v_\nu + 2\theta^2 \bar{\epsilon} \bar{\theta} d + i\theta \sigma^\mu \bar{\epsilon} \bar{\theta}^2 \partial_\mu n \\ &- i\epsilon \sigma^\mu \bar{\theta} \theta \sigma^\nu \bar{\theta} \partial_\mu v_\nu + 2\epsilon \theta \bar{\theta}^2 d - i\epsilon \sigma^\mu \bar{\theta} \theta^2 \bar{\theta} \partial_\mu \bar{\lambda} + i\theta \sigma^\mu \bar{\epsilon} \bar{\theta}^2 \theta \partial_\mu \psi\end{aligned}$$

The details are evidently tedious to verify (and details do not conform with our notation). But the general form is clear with a minimum of computation.

SUSY transformation of components

Reorganize expression into standard expansion of the scalar field in components. $\Rightarrow$ the SUSY transformations in component form:

$$\begin{aligned}
 \delta_\epsilon f &= \epsilon\phi + \bar{\epsilon}\bar{\chi} \\
 \delta_\epsilon\phi_\alpha &= 2\epsilon_\alpha m + \sigma_{\alpha\dot{\beta}}^\mu \bar{\epsilon}^{\dot{\beta}} (i\partial_\mu f + v_\mu) \\
 \delta_\epsilon\bar{\chi}^{\dot{\alpha}} &= 2\bar{\epsilon}^{\dot{\alpha}} n + \epsilon^\beta \sigma_{\beta\dot{\gamma}}^\mu \epsilon^{\dot{\gamma}\dot{\alpha}} [i\partial_\mu f - v_\mu] \\
 \delta_\epsilon m &= \bar{\epsilon}\bar{\lambda} - \frac{i}{2}\partial_\mu\phi\sigma^\mu\bar{\epsilon} \\
 \delta_\epsilon n &= \epsilon\psi + \frac{i}{2}\epsilon\sigma^\mu\partial_\mu\bar{\chi} \\
 \delta_\epsilon v_\mu &= \epsilon\sigma^\mu\bar{\lambda} + \psi\sigma^\mu\bar{\epsilon} + \frac{i}{2}\epsilon\partial_\mu\phi - \frac{i}{2}\partial_\mu\bar{\chi}\epsilon \\
 \delta_\epsilon\bar{\lambda}^{\dot{\alpha}} &= 2\bar{\epsilon}^{\dot{\alpha}} d + \frac{i}{2}\bar{\xi}^{\dot{\alpha}}\partial^\mu v_\mu + i(\epsilon\sigma^\mu\epsilon)^{\dot{\alpha}}\partial_\mu m \\
 \delta_\epsilon\psi_\alpha &= 2\epsilon_\alpha d - \frac{i}{2}\epsilon_\alpha\partial^\mu v_\mu + i(\sigma^\mu\bar{\epsilon})_\alpha\partial_\mu n \\
 \partial_\epsilon d &= \frac{i}{2}\partial_\mu[\psi\sigma^\mu\bar{\epsilon} + \epsilon\sigma^\mu\bar{\lambda}]
 \end{aligned}$$

Important result: the highest component of the superfield (the d -field) transforms to a total derivative ($\theta^2\bar{\theta}^2$ cannot result after ∂_α , $\partial_{\dot{\alpha}}$ derivative, only after ∂_μ derivative.)

Constraints

The general superfield represents the SUSY algebra, but the **representation is reducible**.

Example: the constraints

$$\begin{aligned}\chi(x) &= 0 \\ n(x) &= 0 \\ v_\mu(x) &= i\partial_\mu f(x) \\ \bar{\lambda} &= \frac{i}{2}\partial_\mu \phi(x)\sigma^\mu \\ \psi(x) &= 0 \\ d(x) &= -\frac{i}{4}\square f(x)\end{aligned}$$

are preserved by the SUSY transform.

Remark: the **irreducible** representation defined by these constraints is the **chiral supermultiplet** (in an unfamiliar notation).

The Superspace Derivative

The superspace derivatives

$$\begin{aligned}i\partial_\mu &= P_\mu \\ D_\alpha &= \partial_\alpha - i\sigma_{\alpha\dot{\beta}}^\mu \bar{\theta}^{\dot{\beta}} \partial_\mu \\ \bar{D}_{\dot{\alpha}} &= -\left(\bar{\partial}_{\dot{\alpha}} - i\theta^\beta \sigma_{\beta\dot{\alpha}}^\mu \partial_\mu\right)\end{aligned}$$

are designed such that they **anti-commute** with $Q_\alpha, Q_{\dot{\alpha}}^\dagger$.

They satisfy the fundamental anticommutator with the “wrong” sign

$$\{D_\alpha, \bar{D}_{\dot{\alpha}}\} = -2\sigma_{\alpha\dot{\alpha}}^\mu P_\mu$$

Example: if the components of a general superfields satisfy

$$\bar{D}_{\dot{\alpha}}\Phi = 0$$

then the transformed field will also satisfy this constraint.

This defines the **chiral superfield** in superspace notation.

Constrained Superfields

Define the **superspace coordinate** y^μ such that $\bar{D}_{\dot{\alpha}} y^\mu = 0$:

$$y^\mu = x^\mu - i\theta\sigma^\mu\bar{\theta}$$

Then the general solution to $\bar{D}_{\dot{\alpha}}\Phi = 0$ is a superfield of the form $\Phi(y, \theta)$.

Expand the chiral superfield in components (recalling that squares of Grassmann variables vanish):

$$\begin{aligned}\Phi(y, \theta) &\equiv \phi(y) + \sqrt{2}\theta\psi(y) + \theta^2\mathcal{F}(y) \\ &= \phi(x) - i\theta\sigma^\mu\bar{\theta}\partial_\mu\phi(x) - \frac{1}{4}\theta^2\bar{\theta}^2\partial^2\phi(x) \\ &\quad + \sqrt{2}\theta\psi(x) + \frac{i}{\sqrt{2}}\theta^2\partial_\mu\psi(x)\sigma^\mu\bar{\theta} + \theta^2\mathcal{F}(x)\end{aligned}$$

Remark: the SUSY variation of the **highest component** (the $\mathcal{F}$ -term) must be **a total derivative**.

The kinetic terms in superspace

The kinetic term of a chiral field in superspace formalism:

$$\begin{aligned}
 \int d^4\theta \Phi^\dagger \Phi &= \int d^4\theta \left(\begin{aligned} &\phi^* + i\theta\sigma^\mu\bar{\theta}\partial_\mu\phi^* - \frac{1}{4}\bar{\theta}^2\theta^2\partial^2\phi^* \\ &+ \sqrt{2}\bar{\theta}\psi^\dagger - \frac{i}{\sqrt{2}}\theta\sigma^\mu\partial_\mu\psi^\dagger\bar{\theta}^2 + \bar{\theta}^2\mathcal{F}^* \end{aligned} \right) \\
 &\quad \cdot \left(\begin{aligned} &\phi - i\theta\sigma^\mu\bar{\theta}\partial_\mu\phi - \frac{1}{4}\theta^2\bar{\theta}^2\partial^2\phi \\ &+ \sqrt{2}\theta\psi + \frac{i}{\sqrt{2}}\theta^2\partial_\mu\psi\sigma^\mu\bar{\theta} + \theta^2\mathcal{F} \end{aligned} \right) \\
 &= \mathcal{F}^*\mathcal{F} + \partial^\mu\phi^*\partial_\mu\phi + i\psi^\dagger\bar{\sigma}^\mu\partial_\mu\psi \\
 &\quad - \frac{1}{4}\partial^\mu(\phi^*\partial_\mu\phi + \partial_\mu\phi^*\phi) - \frac{i}{2}\partial_\mu(\psi^\dagger\bar{\sigma}^\mu\psi)
 \end{aligned}$$

Identities:

- $\psi^\dagger\bar{\sigma}^\mu\chi = -\chi\sigma^\mu\psi^\dagger$
- $\theta\sigma^\mu\bar{\theta}\theta\sigma^\nu\bar{\theta} = \frac{1}{2}\eta^{\mu\nu}\theta^2\bar{\theta}^2$

Summary: the superspace integral over the kinetic term of a chiral field:

$$\begin{aligned}
 \int d^4x d^4\theta \Phi^\dagger \Phi &= \int d^4x (\mathcal{F}^*\mathcal{F} + \partial^\mu\phi^*\partial_\mu\phi + i\psi^\dagger\bar{\sigma}^\mu\partial_\mu\psi) \\
 &= \int d^4x \mathcal{L}_{\text{free}}
 \end{aligned}$$

Superpotential

The derivative $\bar{D}_{\dot{\alpha}}$ satisfies the chain rule so the **product of chiral superfields is also a chiral superfield**.

The **superpotential** $W(\Phi)$ is a power series in the chiral superfield, so it is itself a chiral superfield

The SUSY variation of the θ^2 component of a chiral superfield is a total derivative so we can always form the SUSY invariant

$$\int d^4x d^2\theta W(\Phi) = \int d^4x W|_{\theta^2\text{-only}}(\Phi)$$

Component expansion (see next slide for details):

$$\int d^2\theta W(\Phi) = W_a \mathcal{F}^a - \frac{1}{2} W_{ab} \psi^a \psi^b - \partial_\mu \left(\frac{1}{4} W^a \bar{\theta}^2 \partial^\mu \phi_a - \frac{i}{\sqrt{2}} W^a \psi_a \sigma^\mu \bar{\theta} \right)$$

The interaction terms of the WZ-model in superspace notation:

$$\int d^4x d^2\theta W(\Phi) + h.c. = \int d^4x \mathcal{L}_{\text{int}}$$

Details

To work out the component expansion, **first expand y** :

$$W = W_{y=x} + \partial_\mu W|_{y=x} (-i\theta\sigma^\mu\bar{\theta}) + \frac{1}{2} \partial_\mu\partial_\nu W|_{y=x} (-i\theta\sigma^\mu\bar{\theta})(-i\theta\sigma^\nu\bar{\theta})$$

Up to total derivatives we can thus take $y = x$ and immediately find

$$\begin{aligned} W|_{\theta^2-\text{only}} &= [\partial_a W \mathcal{F}^a \theta^2 + \frac{1}{2} \partial_a \partial_b W (\sqrt{2}\theta\psi^a)(\sqrt{2}\theta\psi^b)]_{\theta^2-\text{only}} \\ &= \partial_a W \mathcal{F}^a - \frac{1}{2} \partial_a \partial_b W \psi^a \theta \psi^b \end{aligned}$$

Expanding the derivative terms in θ we find their component form:

$$\begin{aligned} & \partial_\mu [\partial_a W \sqrt{2}\theta\psi^a (-i\theta\sigma^\mu\bar{\theta}) - \frac{1}{4} \partial^\mu W \theta^2 \bar{\theta}^2]_{\theta^2-\text{only}} \\ &= \partial_\mu \left[\frac{i}{\sqrt{2}} \partial_a W (\psi^a \sigma^\mu \bar{\theta}) - \frac{1}{4} W_a \partial^\mu \phi^a \bar{\theta}^2 \right] \end{aligned}$$

Kähler function

More general Lagrangian:

$$\int d^4x d^4\theta K(\Phi^\dagger, \Phi)$$

SUSY invariance: the SUSY variation of the $\theta^2\bar{\theta}^2$ component of **any** superfield is a total derivative.

Terminology: the real function K is called the Kähler function.

It gives general kinetic and other (in general non-renormalizable) interactions.

The Real Superfield

Reality is another constraint that can be consistently imposed on a general scalar superfield:

$$V(x, \theta, \bar{\theta}) = V^\dagger(x, \theta, \bar{\theta})$$

Parametrize the real superfield conveniently

$$\begin{aligned} V(x, \theta, \bar{\theta}) &= C(x) + \theta\chi(x) + \bar{\theta}\chi^\dagger(x) + \frac{1}{2}\theta^2[M(x) + iN(x)] \\ &+ \frac{1}{2}\bar{\theta}^2[M(x) - iN(x)] + \theta\sigma^\mu\bar{\theta}A_\mu(x) + \theta^2[\lambda^\dagger(x) + \frac{1}{2}\partial_\mu\chi(x)\sigma^\mu]\bar{\theta} \\ &+ \bar{\theta}^2\theta[\lambda(x) + \frac{1}{2}\sigma^\mu\partial_\mu\chi^\dagger(x)] + \frac{1}{2}\theta^2\bar{\theta}^2[D(x) - \frac{1}{2}\square C(x)] \end{aligned}$$

Example: real superfield constructed from a chiral superfield Λ

$$\begin{aligned} \Lambda + \Lambda^\dagger &= (A + A^*) + \sqrt{2}\theta\psi + \sqrt{2}\bar{\theta}\psi^\dagger + \theta^2 F + \bar{\theta}^2 F^* - i\theta\sigma^\mu\bar{\theta}\partial_\mu(A - A^*) \\ &+ \frac{1}{\sqrt{2}}\theta^2\partial_\mu\psi\sigma^\mu\bar{\theta} + \frac{1}{\sqrt{2}}\bar{\theta}^2\theta\sigma^\mu\partial_\mu\psi^\dagger - \frac{1}{4}\theta^2\bar{\theta}^2\square(A + A^*) \end{aligned}$$

Extended gauge invariance

Extended gauge invariance acts on the entire real superfield:

$$V \rightarrow V + \Lambda + \Lambda^\dagger$$

It acts as ordinary gauge invariance on the vector field component A_μ :

$$A_\mu \rightarrow A_\mu - i\partial_\mu(A - A^*)$$

A SUSY theory invariant under ordinary gauge symmetry is **automatically invariant under the full extended gauge symmetry**, where the gauge parameter is a chiral superfield, Λ .

The WZ gauge

The WZ-gauge: choose gauge functions of **extended gauge invariance** so three scalar fields $C(x), M(x), N(x)$ and the Weyl spinors $\chi(x), \chi^\dagger(x)$ are transformed to zero.

The vector supermultiplet in the Wess-Zumino gauge:

$$V_{\text{WZ}}^a = \theta \sigma^\mu \bar{\theta} A_\mu^a + \theta^2 \bar{\theta} \lambda^{\dagger a} + \bar{\theta}^2 \theta \lambda^a + \frac{1}{2} \theta^2 \bar{\theta}^2 D^a ,$$

Remark: adjoint indices appropriate for non-abelian generalization. The transformation to WZ-gauge is (much) more complicated in that case, but the final result is the same.

Non abelian gauge invariance

Extended gauge transformation:

$$\exp(T^a V^a) \rightarrow \exp(T^a \Lambda^{a\dagger}) \exp(T^a V^a) \exp(T^a \Lambda^a)$$

Linear approximation:

$$V^a \rightarrow V^a + \Lambda^a + \Lambda^{a\dagger} + \mathcal{O}(V^a \Lambda^a)$$

Higher orders simplify in Wess–Zumino gauge:

$$\begin{aligned} V^a V^b &= \frac{1}{2} \theta^2 \bar{\theta}^2 A^{a\mu} A_{\mu}^b \\ V^a V^b V^c &= 0 \end{aligned}$$

“Field strength” Chiral Superfield

Goal: **make extended gauge symmetry manifest**, and see that $C(x)$, $M(x)$, $C(x)$ and $\chi(x)$, $\chi^\dagger(x)$ are unphysical (they decouple).

Starting from a vector superfield, define the **spinorial** superfield

$$T^a W_\alpha^a = -\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} e^{-T^a V^a} D_\alpha e^{T^a V^a}$$

Expand in components:

$$W_\alpha^a = \lambda_\alpha^a(y) + \theta_\alpha D^a(y) - (\sigma^{\mu\nu} \theta)_\alpha F_{\mu\nu}^a(y) + \theta^2 i \sigma^\mu D_\mu \lambda^{\dagger a}(y),$$

where

$$\sigma^{\mu\nu} = \frac{i}{4} (\sigma^\mu \bar{\sigma}^\nu - \sigma^\nu \bar{\sigma}^\mu)$$

Terminology: W_α is the **“field strength” chiral superfield**.

Interpretation: W_α isolates the physical degrees of freedom from the vector superfield (off-shell, the D -term is included).

Why is W_α “chiral”?

Recall: chiral **scalar** superfields are general superfields satisfying

$$\bar{D}_{\dot{\alpha}}\Phi = 0$$

Definition of field strength superfield:

$$T^a W_\alpha^a = -\frac{1}{4} \bar{D}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} e^{-T^a V^a} D_\alpha e^{T^a V^a}$$

There are just two distinct super-derivatives $\bar{D}_{\dot{\alpha}}$ ($\dot{\alpha} = 1$ and $\dot{\alpha} = 2$) and they “all” anti-commute; so

$$\bar{D}_{\dot{\alpha}} W_\alpha^a = 0$$

Conclusion: W_α^a is **chiral**.

Corollary: the chiral superfield W_α^a is a function of the superspace coordinate $y^\mu = x^\mu - i\theta\sigma^\mu\bar{\theta}$ and θ , **but not $\bar{\theta}$** .

(Proof: these are the quantities annihilated by $\bar{D}_{\dot{\alpha}}$.)

Component Expansion: SUSY QED

Goal: work out the field strength superfield in components.

Definition:

$$W_\alpha = -\frac{1}{4}\bar{D}_{\dot{\alpha}}\bar{D}^{\dot{\alpha}}(D_\alpha V) = D_\alpha V|_{\bar{\theta}^2 \text{ only}}$$

where the vector field (in WZ-gauge):

$$V_{\text{WZ}} = \theta\sigma^\mu\bar{\theta}A_\mu + \theta^2\bar{\theta}\lambda^\dagger + \bar{\theta}^2\theta\lambda + \frac{1}{2}\theta^2\bar{\theta}^2 D ,$$

and the superspace derivative (with the y^μ coordinate kept fixed):

$$D_\alpha = \frac{\partial}{\partial\theta^\alpha} - 2i(\sigma^\mu\bar{\theta})_\alpha \frac{\partial}{\partial y^\mu}$$

Manipulations (details below):

$$\begin{aligned} W_\alpha &= \lambda_\alpha + \theta_\alpha D - 2i(\sigma^\mu\bar{\theta})_\alpha [(\theta\sigma^\nu\bar{\theta}\partial_\mu A_\nu + \theta^2\bar{\theta}\partial_\mu\lambda^\dagger)]_{\bar{\theta}^2 \text{ only}} \\ &= \lambda_\alpha + \theta_\alpha D - (\sigma^{\mu\nu}\theta)_\alpha F_{\mu\nu} + i\theta^2(\sigma^\mu\partial_\mu\lambda^\dagger)_\alpha \end{aligned}$$

Reordering:

$$(\sigma^\mu \bar{\theta})_\alpha \theta \sigma^\nu \bar{\theta} = -\sigma^\mu_{\alpha\dot{\alpha}} \bar{\theta}^{\dot{\alpha}} (\bar{\theta}_{\dot{\beta}} \bar{\sigma}^{\nu\dot{\beta}\beta} \theta_\beta) = \frac{1}{2} (\sigma^\mu \bar{\sigma}^\nu \theta)_\alpha \bar{\theta}^2$$

where

$$\sigma^{\mu\nu} = \frac{i}{4} (\sigma^\mu \bar{\sigma}^\nu - \sigma^\nu \bar{\sigma}^\mu)$$

Some identities:

$$\begin{aligned} \psi \sigma^\mu \chi^\dagger &= -\chi^\dagger \bar{\sigma}^\mu \psi \\ \bar{\theta}^{\dot{\alpha}} \bar{\theta}_{\dot{\beta}} &= -\frac{1}{2} \bar{\theta}^2 \delta_{\dot{\beta}}^{\dot{\alpha}} \\ (\psi \theta)(\chi \theta) &= -\frac{1}{2} (\psi \chi) \theta^2 \end{aligned}$$

Each component field depends on y^μ , the entire W_α depends on y^μ, θ :

$$W_\alpha(y, \theta) = \lambda_\alpha(y) + \theta_\alpha D(y) - (\sigma^{\mu\nu} \theta)_\alpha F_{\mu\nu}(y) + i\theta^2 (\sigma^\mu \partial_\mu \lambda^\dagger(y))_\alpha$$

SUSY Yang–Mills action

Lagrangian in superspace notation:

$$\int d^4x \mathcal{L}_{SYM} = \frac{1}{4} \int d^4x d^2\theta W^{a\alpha} W_\alpha^a + h.c.$$

where

$$\mathcal{L}_{SYM} = -\frac{1}{4} F_{\mu\nu}^a F^{\mu\nu a} + i\lambda^{\dagger a} \bar{\sigma}^\mu D_\mu \lambda^a + \frac{1}{2} D^a D^a$$

An alternate form, as an integral over **all** of superspace:

$$\int d^4x \mathcal{L}_{SYM} = \frac{1}{2} \int d^4x d^4\theta \text{Tr} T^a W^{a\alpha} e^{-T^a V^a} D_\alpha e^{T^a V^a} + h.c.$$

The equivalence:

$$\begin{aligned} T^a W^{a\alpha} &= -\frac{1}{4} \bar{D}^2 (e^{-T^a V^a} D_\alpha e^{T^a V^a}) \\ &= [e^{-T^a V^a} D_\alpha e^{T^a V^a}]_{\theta^2 \text{ only}} \\ &= \int d^2\bar{\theta} [e^{-T^a V^a} D_\alpha e^{T^a V^a}] \end{aligned}$$

Matter Couplings

The chiral superfield Φ transforms under the extended gauge invariance as

$$\Phi \rightarrow e^{-gT^a\Lambda^a}\Phi$$

Standard gauge invariant kinetic terms:

$$\int d^4\theta \Phi^\dagger e^{gT^a V^a} \Phi$$

More generally (to include non-renormalizable interactions):

$$\int d^4\theta K(\Phi^\dagger, e^{gT^a V^a} \Phi)$$