


# Shedding Light on Dark Energy with the SuperNova/Acceleration Probe (SNAP)

Wolfgang Lorenzon  
University of Michigan


Physics Seminar  
Caltech

March 4, 2005



Supernova 1994D

# Hubble's Great Discovery - The Universe is Expanding!



velocity  $\propto$  distance

$$v = H_0 r$$

- In 1929 Hubble measured the redshift (velocity) of nearby galaxies and found that nearly all were moving away from us.
- He used Cepheid variables as “standard candles” to measure distances.
- **Result: The faster they are moving, the farther away they are.**

# Geometry versus Destiny


- General Relativity: Geometry  $\Rightarrow$  Destiny
- For a universe made entirely of matter (with  $\rho \geq 0$ ,  $p = 0$ ) the geometry is determined by “Omega”  $\Omega = \rho_{\text{TOT}}/\rho_{\text{crit}}$ , the ratio between total density of matter and the critical density.

$$\rho_{\text{crit}} = 3H_0^2/8\pi G = 10^{-29} \text{ g/cm}^3 \text{ for } H_0 = 71 \text{ km/s/Mpc (7 H atoms/m}^3\text{)}$$

$\Omega > 1$  Positive curvature, closed universe  $\Rightarrow$  eventual collapse.

$\Omega = 1$  Flat space, open infinite universe  $\Rightarrow$  decelerates to rest.

$\Omega < 1$  Negative curvature, open infinite universe, expands forever.



- Vacuum energy (with  $\rho \geq 0$ ,  $p < 0$ ), such as a cosmological constant, increases in total energy as the universe expands. This will accelerate the expansion of the universe.

# Problems with the Big Bang

## ➤ The Horizon Problem


- CMBR measurements show that the temperature is the same in all directions.
- If we follow the Hubble expansion back in time, regions of the sky separated by more than a few degrees could have never been in contact with each other and could not have come into thermal equilibrium.

## ➤ The Flatness Problem

- If  $\Omega \neq 1$ , it rapidly evolves away from  $\Omega = 1$  as the universe expands.
- If  $\Omega$  is close to unity now, it had to be within  $10^{-28}$  of unity (0.99999999999999999999999999999999) near the beginning of time.
- Only if  $\Omega$  is precisely = 1 would it remain = 1.
- What mechanism could fine-tune the universe to this degree?


# Inflation

- In 1979 Alan Guth proposed an inflationary universe to solve the “horizon problem” and to explain the apparent “flatness” of the universe.
- From  $\sim 10^{-35}$  -  $10^{-33}$  s, the inflationary universe doubled in size every  $10^{-35}$  s, expanding by a factor of  $10^{28}$  and setting  $\Omega = 1$ .
- Quantum fluctuations in the early universe were expanded into the “seeds” that nucleated the large-scale structures in the universe today.
- Inflation was powered by a time-dependent vacuum energy ( $p < 0$ ) that transformed itself into all the matter and energy in the universe today.



# A Startling Discovery



- (1998) **Supernova Cosmology Project** and **High-Z Supernova team** construct a Hubble diagram using Type Ia supernovae looking back 7 Byr (1/2 the age of the universe).
- Both found the **expansion of the universe is accelerating!**



# A Revolution in Cosmology

## Big Bang Nucleosynthesis

## Inflation




Flat universe

$$\Omega_{total} = 1.02 \pm 0.02$$

Baryon Density

$$\Omega_B = 0.044 \pm 0.004$$



- Weak lensing mass census
- Large scale structure measurements

$$\Omega_M = 0.3$$

**New Standard Cosmology:**  
 73±4% Dark Energy  
 27±4% Matter  
 0.5% Bright Stars  
 Matter:  
 22% CDM, 4.4% Baryons,  
 0.3% ν's

# Who Ordered That?!

What's wrong with a Vacuum Energy/Cosmological constant?

- *Why so small?*

Might expect  $\frac{\Lambda}{8\pi G} \sim m_{\text{Planck}}^4$

This is off by  $\sim 120$  orders of magnitude!

Some remarkable unknown symmetry of nature must have cancelled this vacuum energy to allow our universe to spring into existence. But how could it do so and leave a part in  $10^{-120}$  remaining?

- *Why now? Why, why, why?*

$$\frac{\ddot{R}}{R} = -\frac{4\pi G}{3}(\rho + 3p)$$

Matter:  $p = 0 \rightarrow \rho \propto R^{-3}$

Vacuum Energy:  $p = -\rho \rightarrow \rho \propto \text{constant}$

**New Physics:** "Dark energy":  
Dynamical scalar fields, "quintessence",...

**General  
Equation of State:**

$$p = w\rho \rightarrow \rho \propto R^{-3(1+w)}$$

and  $w$  can vary with time

*time*



# Implications of Cosmic Acceleration

There is a previously unseen “dark energy” pervading all of space that is now **accelerating** the expansion of the universe.

Whatever its origin (Cosmological Constant, Vacuum Energy...) the dark energy has a negative pressure ( $p < 0$ ).


This dark energy is the major constituent of the universe today (~2/3) and will continue to increase its share as the universe expands.

The expansion is NOT slowing to a halt and then collapsing (i.e., the universe is not “coming to an end”). In the simplest models, it **will expand forever**.

In the not so distant past (> 5 Byr ago) the universe was dominated by matter and was **decelerating**. Larger and larger structures formed as each new scale entered the horizon.

The formation of structure **ended** when dark energy prevailed over matter.

The largest structures in the universes are now being accelerated beyond our horizon - e.g. Virgo Cluster will be leave our horizon in 118 billion years.



# What we don't know

- Precisely how much mass density ( $\Omega_M$ ) and dark energy density ( $\Omega_{DE}$ ) is there?
- How flat is the universe?
- What is the equation of state ( $w = p/\rho$ ) of the universe and **how has it changed in time?**

Lots of theories, little data!


What is the “dark energy?”  
Theorists have proposed a number of possibilities each with its own unique  $w(t)$ :

- Cosmological constant with  $p = -\rho$  and  $w = -1$ .
- “Quintessence” models with time varying  $-0.4 < w < -0.8$
- Supergravity models
- “Cardassian” expansion
- The “big rip”  $w < -1$

...

# Type Ia SNe:

- Type Ia supernovae (SNe Ia) provide a bright “standard candle” that can be used to construct a Hubble diagram looking back over the last 2/3 of the age of the universe.
- Accretion sends mass of white dwarf star to Chandrasekhar limit leading to gravitational core collapse and a thermonuclear explosion of its outer layers.
- Each one is a strikingly similar explosion event with nearly the **same peak intensity**.




Can measure both intensity and spectra as the supernova brightens and fades over many days.


Comparison of SN Ia redshifts and magnitudes provides straightforward measurement of the changing rate of expansion of the universe:

- **Apparent magnitude measures distance** (time back to explosion)
- **Redshift measures** the total relative **expansion** of the universe since that time
- ☑ Analysis of the spectra characterizes the details of the explosion and helps to control potential systematic errors.


# “Standard” Candles



- Nearby supernovae used to study SNe light curve ( $z < 0.1$ )
- Brightness not quite standard
- Intrinsically brighter SNe last longer
- Correction factor needed





# The Expansion History of the Universe



# It's a SNAP!

To determine what the dark energy is will require a “next generation” experiment, the

## SuperNova/Acceleration Probe



A large wide-field telescope with a Giga-pixel visible/NIR camera and a visible/NIR spectrograph will provide:

- a much larger sample of supernovae (thousands).
- much better control of systematic errors (1 – 2%).
- a much larger range of redshifts (out to  $z = 1.7$ ) to see all the way back through the acceleration and deceleration epochs of the universe.




# SNAP Collaboration



LBLN

G. Aldering, C. Bebek, W. Carithers, C. Day, R. DiGennaro, S. Deustua<sup>†</sup>, D. Groom, M. Hoff, S. Holland, D. Huterer<sup>†</sup>, A. Karcher, A. Kim, W. Kolbe, W. Kramer, B. Krieger, G. Kushner, N. Kuznetsova, R. Lafever, J. Lamoureux, M. Levi, E. Linder, S. Loken, R. Miquel, P. Nugent, H. Oluseyi<sup>†</sup>, N. Palaio, S. Perlmutter, N. Roe, A. Spadafora, H. Von Der Lippe, J-P. Walder, G. Wang



UC Berkeley

M. Bester, E. Commins, G. Goldhaber, H. Heetderks, P. Jelinsky, M. Lampton, D. Pankow, M. Sholl, G. Smoot



Caltech

R. Ellis, R. Massey<sup>†</sup>, A. Refregier<sup>†</sup>, J. Rhodes, R. Smith, K. Taylor



Fermi National Laboratory

J. Annis, F. DeJongh, S. Dodelson, T. Diehl, J. Frieman, L. Hui, S. Kent, P. Limon, J. Marriner, H. Lin, J. Peoples, V. Scarpine, A. Stebbins, C. Stoughton, D. Tucker, W. Wester



Indiana University

C. Bower, N. Mostek, J. Musser, S. Mufson



IN2P3 (France)

P. Astier, E. Barrelet, A. Bonissent, A. Ealet, D. Fouchez<sup>†</sup>, R. Pain, G. Smadja, A. Tilquin, D. Vincent




LAM (France)

S. Basa, R. Malina, A. Mazure, E. Prieto



University of Michigan

B. Bigelow, M. Brown, M. Campbell, D. Gerdes, W. Lorenzon, T. McKay, S. McKee, M. Schubnell, G. Tarlé, A. Tomasch



University of Pennsylvania

G. Bernstein, L. Gladney, B. Jain, D. Rusin



University of Stockholm

R. Amanullah, L. Bergström, A. Goobar, E. Mörtzell

SLAC / Stanford

W. Althouse, R. Blandford, W. Craig, S. Kahn, M. Huffer, P. Marshall



STScI

R. Bohlin, A. Fruchter

Yale University


C. Baltay, W. Emmet, J. Snyder, A. Szymkowiak, D. Rabinowitz, N. Morgan




<sup>†</sup>Institutional affiliation



# Instrument Concept




# Instrument Concept




D=56.6 cm (13.0 mrad)  
0.7 square degrees!


# Focal plane



## Fixed filters atop the sensors





## Integral Field Spectrograph





# Special “Red-Hot” Visible CCDs for SNAP



- New kind of Charged Coupled Device (CCD) developed at Lawrence Berkeley National Lab.
- Better overall response than more costly “thinned” devices in use.
- High-purity “radiation detector” silicon has better radiation tolerance for space applications.
- The CCD’s can be abutted on all four sides enabling very large mosaic arrays.



LBNL “Red Hots”: NOAO September 2001 newsletter


# Near Infrared (NIR) detectors for SNAP

- Large format NIR detectors are a relatively new technology.
- Rockwell Science Center and Raytheon Vision Systems are developing 4 Mpixel (2048 x 2048) NIR focal plane arrays made from HgCdTe (MerCadTel) diodes.
- $\text{Hg}_{1-x}\text{Cd}_x\text{Te}$  composition can be tuned to not “see” long wavelength IR light beyond  $1.7 \mu\text{m}$ , permitting operation at 140K. No active cooling!
- Quantum efficiency, read noise, dark current, uniformity, stability are all being tailored to SNAP requirements.
- Read out by CMOS substrate (> 12 Million transistors) In bump-bonded to HgCdTe diode array (> 4 Million connections).





# Photometry and Spectroscopy Illustration

## Lightcurve & Peak Brightness



## Redshift & SN Properties




$\Omega_M$  and  $\Omega_\Lambda$   
Dark Energy  
Properties

Data




Analysis



Physics

# Launch early in the next decade

1600 kg satellite can be lifted by a Delta IV [recent first flight] to our orbit with margin. Can use equivalent Delta IV, Atlas, or Sea Launch.





# Launch early in the next decade

1600 kg satellite can be lifted by a Delta IV [recent first flight] to our orbit with margin. Can use equivalent Delta IV, Atlas, or Sea Launch.






# Observing

## Step 'n Stare – All Supernovae in all colors

- Repetitive imaging program (SN discovery and light curve measurement)
  - Observe 15 square degrees every three days in all filters “mowing the sky”.
- ~50% of time devoted to spectroscopy of individual SNe near maximum light.





# Data Sheets to Cosmological Parameters


Lightcurve & Peak Brightness



Redshift & SN Properties



Data Sheets for each SN





Instrument




Observed Data

Mission Plan


# Data Sheets to Cosmological Parameters




$\Omega_M$  and  $\Omega_\Lambda$   
Dark Energy Properties  
( $w_0$  and  $w'$ )



Data



Analysis




Physics


Calibration, External SN Obs.

Priors, External Cosmology

# Simulated SNAP Data



# Understanding Dark Energy



based on  
Weller & Albrecht (2001)

# Determination of Cosmological Parameters

**SNAP** will measure

(if  $w = -1$ )

$\Omega_\Lambda$  to  $\pm 0.02$  and


$\Omega_M$  to  $\pm 0.03$ .

With prior  $\Omega_M$  measured by other techniques (e.g. CMB) to  $\pm 0.03$ ,

**SNAP** will measure

$\Omega_M$  to  $\pm 0.01$  and


$w$  to  $\pm 0.05$ .



# Dark Energy Equation of State


Is  $w = -1$ ? By measuring the evolution of  $w = p/\rho$  to high precision  
SNAP will determine the nature of the Dark Energy.

For a flat universe  
with prior  $\Omega_M$   
measured  
to  $\pm 0.03$ ,  
SNAP will measure  
 $w_0$  to  $\pm 0.05$  and  
 $w'$  to  $\pm 0.27$ .




# Why go to high redshifts?

- Dark energy can be detected at low redshift (SCP, High-z). To determine **what it is** and not just **that it is**, requires measurements over both the acceleration and deceleration epochs.
- This long reach breaks essential degeneracies which low redshift data alone cannot.




## SNAP will

- ✓ probe the variability of  $w$ , providing an essential clue to the nature of DE.
- ✓ measure  $w_0$  precisely to determine whether it is a cosmological constant.






# Rest frame B and V shift to NIR



Simulated SNAP observations of high redshift SNe

# This can't be done on the ground!





Simulated 8m telescope ground based observations of high redshift SNe

# NIR available only in space

Crucial near-infrared observations are impossible from the ground

- Sky is very bright in NIR, about 500x brighter at  $1.5\mu\text{m}$ , like observing the sky in Manhattan
- Sky is not transparent in NIR, absorption due to  $\text{H}_2\text{O}$  molecular absorption bands is very strong and extremely variable



# Confronting Systematic Errors

Spectroscopy fixes these

evolution

- Shifting distribution of progenitor mass/metallicity/C-O
- Shifting distribution of SN physics parameters:
  - Amount of Nickel fused in explosion
  - Distribution of Nickel
  - Kinetic energy of explosion
  - Opacity of atmosphere's inner layers
  - Metallicity

NIR fixes these

- Gravitational Lensing (de)amplification

## • Dust/Extinction

- Dust that reddens
- Evolving gray dust
  - Clumpy
  - Homogeneous
- Galactic extinction model

## • Observational biases

- Malmquist bias differences
- non-SN Ia contamination
- K-correction uncertainty
- Color zero-point calibration


# Beyond Dark Energy

1. **Galaxy studies:** evolution and clustering.
2. **Galaxy clusters:** identification of high redshift galaxy clusters, faint and small constituents
3. **High-z quasar studies:** mapping quasars to  $z = 10$ , probing the re-ionization of the universe.
4. **Transients;** Gamma Ray Bursts, Quasars/Active Galactic Nuclei, outer solar system objects
5. **Cool stars in the Milky Way**
6. **Lensing:** evolution of the galaxy/dark matter correlation, determination of galaxy cluster masses, etc.
7. **Targets:** Identification of targets for the James Webb Space Telescope (JWST), California Extremely Large Telescope (CELT), etc...

# SNAP Observation Program

**Base SNAP survey:  
15 square degrees  
near ecliptic poles**

- **~9,000 × as large as Hubble Deep Field, same resolution but 1.5 mag deeper (in nine optical and IR bands)**



# Conclusions

Dark energy is the dominant fundamental constituent of our Universe, yet we know very little about it.

SNAP will show how the expansion rate has varied over the history of the Universe and test theories of dark energy.

A vigorous R&D program, supported by the DoE is underway, leading to an expected launch early in the next decade.

NASA and DoE have agreed to partner on a Joint Dark Energy Mission (JDEM). SNAP is a prime candidate for JDEM.

**THE END**



## The FY 2005 Budget

"The control level is at the High Energy Physics level. The conferees encourage the Department to **proceed** with the **Dark Energy Mission** even if the primary science of the mission and mission development must be pursued by the Department so as to avoid schedule delays resulting from implementing the mission jointly with NASA. **International** cooperation and **appropriate launch** arrangements should be **pursued** where appropriate. The conferees recognize that an excellent and energized science team has been assembled for this exciting mission"