


Complex Systems 535/Physics 508: Homework 3


1. One can calculate the diameter of certain types of networks exactly:

- (i) What is the diameter of a clique?
- (ii) What is the diameter of a square portion of square lattice, with L edges (or equivalently $L + 1$ vertices) along each side, like this:



What is the diameter of the corresponding hypercubic lattice in d dimensions with L edges along each side? Hence what is the diameter of such a lattice as a function of the number n of vertices?

- (iii) A Cayley tree is a symmetric regular tree in which each vertex is connected to the same number k of others, until we get out to the leaves, like this:



(We have $k = 3$ in this picture.)

Show that the number of vertices reachable in ℓ steps from the central vertex is $k(k - 1)^{\ell - 1}$. Hence find an expression for the diameter of the network in terms of k and the number of vertices n .

- (iv) Which of the networks in parts (i), (ii), and (iii) displays the small-world effect, defined as having a diameter that increases as $\log n$ or slower?
2. Suppose a network has a degree distribution that follows the exponential form $p_k = Ce^{-\lambda k}$, where C and λ are constants.
- (i) Find C as a function of λ .
 - (ii) Calculate the fraction P of vertices that have degrees greater than or equal to k .
 - (iii) Calculate the fraction W of ends of edges that are attached to vertices of degree greater than or equal to k .

- (iv) Hence show that for this degree distribution the Lorenz curve—the equivalent of Eq. (8.23) in the course-pack—is given by

$$W = P + \frac{1 - e^{-\lambda}}{\lambda} P \ln P.$$

- (v) What is the equivalent of the “80–20” rule for such a network with $\lambda = 1$? That is, what fraction of the “richest” nodes in the network have 80% of the “wealth”?
3. A particular network is believed to have a degree distribution that follows a power law. A random sample of vertices is taken and their degrees measured. The degrees of the first twenty vertices with degrees 10 or greater are:

16	17	10	26	13
14	28	45	10	12
12	10	136	16	25
36	12	14	22	10

Estimate the exponent α of the power law and the error on that estimate.

4. Consider the following simple and rather unrealistic model of a network: each of n vertices belongs to one of g groups. The m th group has n_m vertices and each vertex in that group is connected to others in the group with independent probability $p_m = A(n_m - 1)^{-\beta}$, where A and β are constants, but not to any vertices in other groups. Thus this network takes the form of a set of disjoint groups or communities.
- (i) Calculate the expected degree $\langle k \rangle$ of a vertex in group m .
 - (ii) Calculate the expected value \bar{C}_m of the local clustering coefficient for vertices in group m .
 - (iii) Hence show that $\bar{C}_m \propto \langle k \rangle^{-\beta/(1-\beta)}$. What value would β have to have for the expected value of the local clustering to fall off as $\langle k \rangle^{-3/4}$, as has been conjectured by some researchers?