

## PROBLEM SET 7 (DUE ON THURSDAY, APRIL 8)

(All Exercises are references to the November 18, 2017 version of *Foundations of Algebraic Geometry* by R. Vakil.)

- Problem 1.** Exercise 19.11.B (a map of projective varieties not arising from a map of (regraded) graded rings)
- Problem 2.** Suppose  $(E, p)$  is a (smooth) elliptic curve in Weierstrass form  $y^2 = x^3 + ax + b$  (along with the point at infinity  $p$ , so really  $E = \text{Proj } k[x, y, z]/(y^2z - x^3 - axz^2 - bz^3)$ ). Show that the rational section  $dx/y$  of  $\Omega_{E/k}$  actually has no zeroes or poles in  $E$ , so it generates  $\Omega_{E/k} \cong \mathcal{O}_E$ .
- Problem 3.** Let  $\iota : C \rightarrow \mathbb{P}_k^2$  be the inclusion of a smooth plane curve of degree  $d > 0$ . Let  $\mathcal{F} = \iota^* \Omega_{\mathbb{P}^2/k}$ , a rank 2 vector bundle on  $C$ . Compute the Euler characteristic  $\chi(C, \mathcal{F})$ .
- Problem 4.** Exercise 21.5.D ( $h^0(X, K_X)$  for a degree  $d$  surface  $X$  in  $\mathbb{P}^3$ . Hint: use adjunction to compute  $K_X$  as the restriction of a line bundle on  $\mathbb{P}^3$ , then pushforward to  $\mathbb{P}^3$  and compute cohomology there using a long exact sequence and the known values of  $h^i(\mathbb{P}^3, \mathcal{O}(m))$ .)