

2. For generalized Jeffrey Fluid:

$$\underline{\tau} + \lambda_1 \left[\frac{\Delta}{\Delta} - a(\underline{\tau} \cdot \underline{\dot{\gamma}} + \underline{\dot{\gamma}} \cdot \underline{\tau}) \right] = 2\eta_0 \left[\underline{\dot{\gamma}} + \lambda_2 \underline{\dot{\gamma}} \right]$$

where: $\underline{\Delta} = \frac{D(\underline{\tau})}{Dt} + \underline{\omega} \cdot (\underline{\tau}) - (\underline{\tau}) \cdot \underline{\omega}$ $\nearrow = \begin{pmatrix} \frac{\partial \tau_{11}}{\partial t} & \frac{\partial \tau_{12}}{\partial t} \\ \frac{\partial \tau_{21}}{\partial t} & \frac{\partial \tau_{22}}{\partial t} \\ \vdots & \vdots \end{pmatrix}$

For shear flow $\begin{cases} u = \dot{\gamma} y \\ v = 0 \end{cases} \quad \nabla u = \begin{pmatrix} 0 & \dot{\gamma} & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \dot{\gamma}$

$$\underline{\dot{\gamma}} = \frac{1}{2} (\nabla u + \nabla u^T) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \frac{\dot{\gamma}}{2} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \alpha$$

$$\underline{\dot{\omega}} = \frac{1}{2} (\nabla u - \nabla u^T) = \begin{pmatrix} 0 & -1 & 0 \\ +1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \frac{\dot{\gamma}}{2} = \begin{pmatrix} 0 & -1 & 0 \\ +1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \alpha$$

Subs: $\xrightarrow{\text{steadily}} \xrightarrow{\text{fully developed}}$

$$\tau_{ij} + \lambda_1 \left[\dot{\tau}_{ij} + v_k \tau_{ij,k} + \omega_{ik} \tau_{kj} - \tau_{ik} \omega_{kj} - a(\tau_{ik} \dot{\gamma}_{kj} + \dot{\gamma}_{ik} \tau_{kj}) \right] = 2\eta_0 \left[\dot{\gamma}_{ij} + \lambda_2 \left(\frac{D \dot{\gamma}_{ij}}{Dt} + \omega_{ik} \dot{\gamma}_{kj} - \dot{\gamma}_{ik} \omega_{kj} \right) \right]$$

note $\tau_{12} = \tau_{21}$, & $\tau_{13} = \tau_{23} = \tau_{33} = 0$

the non-trivial terms are:

$$\tau_{11} + \lambda_1 \left(-\alpha \tau_{21} + \alpha \tau_{12} - a(\alpha \tau_{12} + \alpha \tau_{21}) \right) = 2\eta_0 \left(0 + \lambda_2 (\alpha^2 + \alpha^2) \right)$$

$-2\alpha \tau_{12} - 2a\alpha \tau_{12}$

$$\tau_{11} + 2\lambda_1 \alpha \tau_{12} (1+a) = -4\eta_0 \lambda_2 \alpha^2$$

$$\textcircled{1} \dots \tau_{11} + \lambda_1 \dot{\gamma} \tau_{12} (1+a) = -\eta_0 \lambda_2 \dot{\gamma}^2$$

$$\tau_{12} + \lambda_1 \left[-\alpha \tau_{22} - \alpha \tau_{11} - a(\alpha \tau_{11} + \alpha \tau_{22}) \right] = 2\eta_0 \left[\alpha + \lambda_2 (0 - 0) \right]$$

$$\textcircled{2} \dots \tau_{12} + \lambda_1 \left[\alpha \tau_{11} (1-a) - \alpha \tau_{22} (1+a) \right] = 2\eta_0 \alpha$$

$$\tau_{22} + \lambda_1 \left[+\tau_{22} \alpha + \tau_{11} \alpha - a(\tau_{11} \alpha + \tau_{22} \alpha) \right] = 2\eta_0 \left[0 + \lambda_2 (+\alpha^2 + \alpha^2) \right]$$

$$\textcircled{3} \dots \tau_{22} + \lambda_1 \dot{\gamma} \tau_{12} (1-a) = +\eta_0 \lambda_2 \dot{\gamma}^2$$

Subs ①, ③ into ②

$$\tau_{12} + (\lambda_1 \dot{\gamma} \tau_{12} (1-a) - \eta_0 \lambda_2 \dot{\gamma}^2) \frac{\lambda_1 \dot{\gamma} (1-a)}{2} - \frac{\lambda_1 \dot{\gamma} (1-a)}{2} (-\lambda_1 \dot{\gamma} \tau_{12} (1-a) + \eta_0 \lambda_2 \dot{\gamma}^2) \\ = \cancel{\eta_0 \lambda_2 \dot{\gamma}^2} = \eta_0 \dot{\gamma}$$

$$\Rightarrow \tau_{12} \left[1 + \lambda_1^2 \dot{\gamma}^2 (1-a^2) \right] - \eta_0 \lambda_1 \lambda_2 \dot{\gamma}^3 = \cancel{\eta_0 \lambda_2 \dot{\gamma}^2} = \eta_0 \dot{\gamma}$$

$$\Rightarrow \textcircled{a} \tau_{12} = \frac{\eta_0 \dot{\gamma} (1 + \lambda_1 \lambda_2 \dot{\gamma}^2)}{1 + \lambda_1^2 \dot{\gamma}^2 (1-a^2)}$$

$$\textcircled{b} \tau_{11} - \tau_{22} = -2\eta_0 \lambda_2 \dot{\gamma}^2 + 2\lambda_1 \dot{\gamma} \tau_{12}$$

$$= -2\eta_0 \lambda_2 \dot{\gamma}^2 + \frac{2\eta_0 \lambda_1 \dot{\gamma}^2 (1 + \lambda_1 \lambda_2 \dot{\gamma}^2)}{1 + \lambda_1^2 \dot{\gamma}^2 (1-a^2)} \\ = \frac{2\eta_0 \dot{\gamma}^2 (\lambda_1 - \lambda_2) + 2\eta_0 \lambda_1^2 \lambda_2 \dot{\gamma}^4 a^2}{1 + \lambda_1^2 \dot{\gamma}^2 (1-a^2)}$$

$$\textcircled{c} \tau_{22} - \tau_{33} = \eta_0 \lambda_2 \dot{\gamma}^2 - \lambda_1 \dot{\gamma} (1-a) - \frac{\eta_0 \dot{\gamma} (1 + \lambda_1 \lambda_2 \dot{\gamma}^2)}{1 + \lambda_1^2 \dot{\gamma}^2 (1-a^2)}$$

$$= \frac{\eta_0 \dot{\gamma}^2 [\lambda_2 - \lambda_1 (1-a)] + \eta_0 \lambda_1^2 \lambda_2 \dot{\gamma}^4 (a-a^2)}{1 + \lambda_1^2 \dot{\gamma}^2 (1-a^2)}$$