

1 The interior and the closure of a set

Definition 1.1. (Interior of a set.) Let (X, d) be a metric space, and $A \subseteq X$ a subset. Then the *interior of A* , denoted $\text{Int}(A)$ or $\overset{\circ}{A}$, is defined to be the set

$$\text{Int}(A) = \{a \in A \mid a \text{ is an interior point of } A\}.$$

Note that $\text{Int}(A) \subseteq A$. We will see in the exercises that $\text{Int}(A)$ is an open set, and it is in a sense the largest open subset of A .

Definition 1.2. (Closure of a set.) Let (X, d) be a metric space, and $A \subseteq X$ a subset. Then the *closure of A* , denoted \overline{A} , is defined to be the set

$$\overline{A} = \{x \in X \mid \text{for every } r > 0 \text{ the ball } B_r(x) \text{ contains a point of } A\}.$$

We will see that \overline{A} is a closed set, and that in a sense it is the smallest closed set containing A .

Example 1.3. What is the closure of the open set $B_1(0, 0) \subseteq \mathbb{R}^2$?

In-class Exercises

1. For this problem we introduce the following terminology.

Definition 1.4. (Neighbourhood of a point x .) Let (X, d) be a metric space, and $x \in X$. Then any open set U containing x is called an *open neighbourhood of x* , or simply a *neighbourhood of x* .

- (a) Prove the following.

Theorem 1.5. (Equivalent definition of interior point.) For a subset V of a metric space X , a point $x \in V$ is an interior point of V if and only if there exists an open neighbourhood U of x that is contained in V .

- (b) Prove the following.

Theorem 1.6. (Equivalent definition of closure.) For a subset A of a metric space X , the closure of A is equal to the set

$$\overline{A} = \{x \in X \mid \text{every neighbourhood } U \text{ of } x \text{ contains a point of } A\}.$$

2. Prove the following theorem.

Theorem 1.7. Let (X, d) be a metric space, and $A \subseteq X$ a subset.

- | | |
|---|--|
| (i) $\text{Int}(A) \subseteq A$ | (v) $\text{Int}(A)$ is open in X |
| (ii) A is open if and only if $A = \text{Int}(A)$ | (vi) $\text{Int}(A)$ is the largest open subset of A in the following sense: If $U \subseteq A$ is any open subset of A , then $U \subseteq \text{Int}(A)$ |
| (iii) If $A \subseteq B$ then $\text{Int}(A) \subseteq \text{Int}(B)$ | |
| (iv) $\text{Int}(\text{Int}(A)) = \text{Int}(A)$ | |

3. Prove the following theorem.

Theorem 1.8. Let (X, d) be a metric space, and $A \subseteq X$ a subset.

- | | |
|--|--|
| (i) $A \subseteq \bar{A}$ | (v) \bar{A} is closed in X |
| (ii) If $A \subseteq B$ then $\bar{A} \subseteq \bar{B}$ | (vi) \bar{A} is the smallest closed set containing A , in the following sense: If $A \subseteq C$ for some closed set C , then $\bar{A} \subseteq C$ |
| (iii) A is closed if and only if $A = \bar{A}$ | |
| (iv) $\overline{\bar{A}} = \bar{A}$ | |

4. **(Optional).** Let A be a subset of a metric space (X, d) . Explore the relationships between the sets

$$\text{Int}(X \setminus A) \quad X \setminus \text{Int}(A) \quad \overline{X \setminus A} \quad X \setminus \bar{A}$$

Determine which of these sets are necessarily equal or necessarily subsets of one another. Give counterexamples to show where equality or containment fails.

5. **(Optional).** Let $A_i, i \in I$, be a collection of subsets of a metric space (X, d) . For each of the following statements, either prove the statement, or construct a counterexample.

- | | | | |
|---|---|--|--|
| (a) $\text{Int} \left(\bigcup_{i \in I} A_i \right) \subseteq \bigcup_{i \in I} \text{Int}(A_i)$ | (c) $\text{Int} \left(\bigcap_{i \in I} A_i \right) \subseteq \bigcap_{i \in I} \text{Int}(A_i)$ | | |
| (b) $\text{Int} \left(\bigcup_{i \in I} A_i \right) \supseteq \bigcup_{i \in I} \text{Int} A_i$ | (d) $\text{Int} \left(\bigcap_{i \in I} A_i \right) \supseteq \bigcap_{i \in I} \text{Int}(A_i)$ | | |
| (e) $\overline{\bigcup_{i \in I} A_i} \subseteq \bigcup_{i \in I} \bar{A}_i$ | (f) $\overline{\bigcup_{i \in I} A_i} \supseteq \bigcup_{i \in I} \bar{A}_i$ | (g) $\overline{\bigcap_{i \in I} A_i} \subseteq \bigcap_{i \in I} \bar{A}_i$ | (h) $\overline{\bigcap_{i \in I} A_i} \supseteq \bigcap_{i \in I} \bar{A}_i$ |

6. **(Optional).** Prove the following equivalent definition of continuity.

Theorem (An equivalent definition of continuity). Let (X, d_X) and (Y, d_Y) be metric spaces. Then a map $f : X \rightarrow Y$ is continuous if and only if

$$f(\bar{A}) \subseteq \overline{f(A)} \quad \text{for every subset } A \subseteq X.$$

7. **(Optional).** For a metric (X, d) , let $x_0 \in X$ and $r > 0$. You proved on the homework that the set

$$C_r(x_0) = \{x \in X \mid d(x, x_0) \leq r\}$$

is closed. Explain why $C_r(x_0)$ always contains the closure of the ball $B_r(x_0)$. Give an example of a metric space where $C_r(x_0)$ is equal to $\overline{B_r(x_0)}$ for every $r > 0$ and x_0 , and give an example of a metric space and x_0, r such that $C_r(x_0)$ is a strict subset of $\overline{B_r(x_0)}$.